arXiv:1107.2692v1 [math.SP] 13 Jul 2011

ON THE RESONANCES AND EIGENVALUES FOR A 1D HALF-CRYSTAL WITH LOCALISED IMPURITY

EVGENY L. KOROTYAEV AND KARL MICHAEL SCHMIDT

Abstract. We consider the Schrödinger operator H on the half-line with a periodic potential p plus a compactly supported potential q. For generic p, its essential spectrum has an infinite sequence of open gaps. We determine the asymptotics of the resonance counting function and show that, for sufficiently high energy, each non-degenerate gap contains exactly one eigenvalue or antibound state, giving asymptotics for their positions. Conversely, for any potential q and for any sequences $(\sigma_n)_1^\infty, \sigma_n \in \{0,1\}$, and $(\varkappa_n)_1^\infty \in \ell^2, \varkappa_n \geqslant 0$, there exists a potential p such that $\varkappa_n$ is the length of the n-th gap, $n \in \mathbb{N}$, and H has exactly σ_n eigenvalues and $1-\sigma_n$ antibound state in each high-energy gap. Moreover, we show that between any two eigenvalues in a gap, there is an odd number of antibound states, and hence deduce an asymptotic lower bound on the number of antibound states in an adiabatic limit.

1. Introduction and main results

Consider the Schrödinger operator H acting in the Hilbert space $L^2(\mathbb{R}_+)$ and given by

$$H = H_0 + q, \qquad H_0 f = -f'' + pf$$

with the boundary condition $f(0) = 0$. Here p is 1-periodic and q is compactly supported and they satisfy

$$p \in L^1_{real}(\mathbb{R}/\mathbb{Z}), \qquad q \in \mathcal{Q}_t, \tag{1.1}$$

where $\mathcal{Q}_t = \{q \in L^2_{real}(\mathbb{R}_+) \mid \sup(\mathrm{supp}(q)) = t\}$ and we keep $t > 0$ fixed throughout. For later use, we set $n_t = \inf_{n\in\mathbb{N}, n\geqslant t} n$. The spectrum of H_0 consists of an absolutely continuous part $\sigma_{ac}(H_0) = \bigcup\limits_{n\in\mathbb{N}} \mathfrak{S}_n$ plus at most one eigenvalue in each non-empty gap γ_n, $n \in \mathbb{N}$, [E2], [Zh3]), where the bands $\mathfrak{S}_n$ and gaps γ_n are given by (see Fig. 1)

$$\mathfrak{S}_n = [E^+_{n-1}, E^-_n], \qquad \gamma_n = (E^-_n, E^+_n) \quad (n \in \mathbb{N}),$$

and the E_n satisfy

$$E^+_0 < E^-_1 \leqslant E^+_1 \cdots \leqslant E^+_{n-1} < E^-_n \leqslant E^+_n < \dots \tag{1.2}$$

It is known that there are infinitely many non-degenerate gaps, i.e. $E^-_n < E^+_n$, unless p is arbitrarily often differentiable [Ho], and all gaps are non-degenerate generically [MO], [S1]. Without loss of generality, we may assume $E^+_0 = 0$. The sequence ((1.2)) is the spectrum of the equation

$$-y'' + p(x)y = \lambda y \tag{1.3}$$

Saint-Petersburg University; korotyaev@gmail.com
School of Mathematics, Cardiff University. Senghennydd Road, CF24 4AG Cardiff, Wales UK; SchmidtKM@cf.ac.uk

Date: July 15, 2011.
1991 Mathematics Subject Classification. 34A55, (34B24, 47E05).
Key words and phrases. resonances, scattering, periodic potential, S-matrix.



FIGURE 1. The cut domain $\mathbb{C} \setminus \cup \mathfrak{S}_n$ and the cuts (bands) $\mathfrak{S}_n = [E_{n-1}^+, E_n^-], n \geqslant 1$

with the condition of 2-periodicity, $y(x+2) = y(x)$ $(x \in \mathbb{R})$. We also set $\gamma_0 = (-\infty, E_0^+)$. If a gap degenerates, $\gamma_n = \emptyset$ for some n, then the corresponding bands $\mathfrak{S}_n$ and $\mathfrak{S}_{n+1}$ touch. This happens when $E_n^- = E_n^+$; this number is then a double eigenvalue of the 2-periodic problem (1.3). The lowest eigenvalue $E_0^+ = 0$ is always simple and has a 1-periodic eigenfunction. Generally, the eigenfunctions corresponding to eigenvalues $E_{2n}^\pm$ are 1-periodic, those for $E_{2n+1}^\pm$ are 1-anti-periodic in the sense that $y(x+1) = -y(x)$ $(x \in \mathbb{R})$.

Consider the operator $\mathscr{H} y = -y'' + (p+q)y$ on the real line, where p is periodic and and q is compactly supported. The spectrum of the operator $\mathscr{H}$ consists of an absolutely continuous part $\sigma_{ac}(\mathscr{H}) = \sigma_{ac}(H_0)$ plus a finite number of simple eigenvalues in each non-empty gap $\gamma_n, n \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$ ([Rb], [F1]), and has at most two eigenvalues in every open gap γ_n with sufficiently large n ([Rb]). If $q_0 := \int_{\mathbb{R}} q(x)dx \neq 0$, then $\mathscr{H}$ has precisely one eigenvalue ([Zh1], [F2], [GS]) and one antibound state [K4] in each non-empty gap γ_n with sufficiently large n. If $q_0 = 0$, then there are, roughly speaking, either two eigenvalues and no antibound states or no eigenvalues and two antibound states in each non-empty gap γ_n with sufficiently large n ([K4]). Similarly, the spectrum of H defined in (1.1) consists of an absolutely continuous part $\sigma_{ac}(H) = \sigma_{ac}(H_0)$ plus a finite number of simple eigenvalues in each non-empty gap γ_n, $n \in \mathbb{N}_0$. This follows from the corresponding property of $\mathscr{H}$ by the Glazman decomposition principle.

Throughout the paper, we shall denote by $\vartheta(x,z)$, $\varphi(x,z)$ the two solutions forming the canonical fundamental system of the unperturbed equation $-y'' + py = z^2 y$, i.e., satisfying the initial conditions $\varphi'(0,z) = \vartheta(0,z) = 1$ and $\varphi(0,z) = \vartheta'(0,z) = 0$. (Here and in the following $'$ denotes the derivative w.r.t. the first variable.) The Lyapunov function (Hill discriminant) of the periodic equation is then defined by $\Delta(z) = \frac{1}{2}(\varphi'(1,z) + \vartheta(1,z))$. The function $\lambda \mapsto \Delta^2(\sqrt{\lambda})$ is entire, where we take the square root to be positive on the positive real axis and to map the remainder of the complex plane into the upper half-plane. For the function $(1 - \Delta^2(\sqrt{\lambda}))^{\frac{1}{2}}$ $(\lambda \in \overline{\mathbb{C}}_+)$, we fix the branch by the condition $(1 - \Delta^2(\sqrt{\lambda + i0}))^{\frac{1}{2}} > 0$ for $\lambda \in \mathfrak{S}_1 = [E_0^+, E_1^-]$, and introduce the two-sheeted Riemann surface Λ of $(1 - \Delta^2(\sqrt{\lambda}))^{\frac{1}{2}}$ obtained by joining the upper and lower rims of two copies of the cut plane $\mathbb{C} \setminus \sigma_{ac}(H_0)$ in the usual (crosswise) way, see e.g. [F3]. We denote the n-th gap on the first, physical sheet Λ_1 by $\gamma_n^{(1)}$ and its counterpart on the second, nonphysical sheet Λ_2 by $\gamma_n^{(2)}$, and set

$$\gamma_n^c := \overline{\gamma}_n^{(1)} \cup \overline{\gamma}_n^{(2)}. \tag{1.4}$$

It is well known (see e.g. (2.18)) that, for each $h \in C_0^\infty(\mathbb{R}_+), h \neq 0$, the function $f(\lambda) = ((H-\lambda)^{-1}h, h)$ has a meromorphic extension from the physical sheet Λ_1 to the whole Riemann surface Λ. Moreover, if f has a pole at some $\lambda_0 \in \Lambda_1$ for some h, then λ_0 is an eigenvalue (bound state) of H and $\lambda_0 \in \bigcup\limits_{n\in\mathbb{N}_0} \gamma_n^{(1)}$.

Definition. Let $f(\lambda) = ((H-\lambda)^{-1}h, h)$ $(\lambda \in \Lambda)$ for some $h \in C_0^\infty(\mathbb{R}_+), h \neq 0$.
1) If $f(\lambda)$ has a pole at some $\lambda_0 \in \Lambda_2$, $\lambda_0 \neq E_n^\pm, n \geqslant 0$, we call λ_0 a *resonance*.
2) A point $\lambda_0 = E_n^\pm, n \geqslant 0$, is called a *virtual state* if the function $z \mapsto f(\lambda_0 + z^2)$ has a pole at 0.
3) A point $\lambda_0 \in \Lambda$ is called a *state* if it is either a bound state or a resonance or a virtual state. Its *multiplicity* is the multiplicity of the corresponding pole. We denote by $\mathfrak{S}_{st}(H)$ the set of all states. If $\lambda_0 \in \gamma_n^{(2)}, n \geqslant 0$, then we call λ_0 an *antibound state*.

As a telling example we consider the states of the operator H_0 for the case $p \neq \text{const}, q = 0$, see [Zh3], [HKS]. Let $f_0(\lambda) = ((H_0-\lambda)^{-1}h, h)$ for some $h \in C_0^\infty(\mathbb{R}_+)$. It is well known that the function f_0 is meromorphic on the physical sheet Λ_1 and has a meromorphic extension into Λ (see e.g. (2.18)). For each $\gamma_n^c \neq \emptyset, n \geqslant 1$, there is exactly one state $\lambda_n^0 \in \gamma_n^c$ of H_0 and its projection onto the complex plane coincides with the n-th eigenvalue, μ_n^2, of the Dirichlet boundary value problem

$$-y_n'' + py_n = \mu_n^2 y_n, \quad y_n(0) = y_n(1) = 0, \qquad x \in [0,1], \quad n \geqslant 1.$$

Moreover, exactly one of the following three cases holds,
1) $\lambda_n^0 \in \gamma_n^{(1)}$ is an eigenvalue,
2) $\lambda_n^0 \in \gamma_n^{(2)}$ is an antibound state, or
3) $\lambda_n^0 \in \{E_n^+, E_n^-\}$ is a virtual state.

There are no other states of H_0, so H_0 has only eigenvalues, virtual states and antibound states. If there are exactly $N \geqslant 1$ nondegenerate gaps in the spectrum of $\sigma_{ac}(H_0)$, then the operator H_0 has exactly N states; the closed gaps $\gamma_n = \emptyset$ do not contribute any states. In particular, if $\gamma_n = \emptyset$ $(n \geqslant 1)$, then $p = 0$ ([MO], [K5]) and H_0 has no states. A more detailed description of the states of H_0 is given in Lemma 2.1 below.

We shall need the following results from the inverse spectral theory for the unperturbed operator H_0. Defining the mapping $p \mapsto \xi = (\xi_n)_1^\infty$, where the components $\xi_n = (\xi_{1n}, \xi_{2n}) \in \mathbb{R}^2$ are given by

$$\xi_{1n} = \frac{E_n^- + E_n^+}{2} - \mu_n^2, \qquad \xi_{2n} = \left|\frac{|\gamma_n|^2}{4} - \xi_{1n}^2\right|^{\frac12} a_n, \qquad a_n = \begin{cases} +1 & \text{if } \lambda_n^0 \text{ is an eigenvalue,} \\ -1 & \text{if } \lambda_n^0 \text{ is a resonance,} \\ 0 & \text{if } \lambda_n^0 \text{ is a virtual state,} \end{cases}$$

we have the following result from [K5], [K7]:
The mapping $\xi : \mathcal{H} \to \ell^2 \oplus \ell^2$ is a real analytic isomorphism between the real Hilbert spaces $\mathcal{H} = \{p \in L^2(0,1) \mid \int_0^1 p(x)dx = 0\}$ and $\ell^2 \oplus \ell^2$, and the estimates

$$\|p\| \leqslant 4\|\xi\|(1 + \|\xi\|^{\frac13}), \qquad \|\xi\| \leqslant \|p\|(1 + \|p\|)^{\frac13} \tag{1.5}$$

hold, where $\|p\|^2 = \int_0^1 p^2(x)dx$ and $\|\xi\|^2 = \frac14 \sum |\gamma_n|^2$.
Moreover, given any non-negative sequence $\varkappa = (\varkappa_n)_1^\infty \in \ell^2$, there are unique 2-periodic eigenvalues $E_n^\pm$ $(n \in \mathbb{N}_0)$, for some $p \in \mathcal{H}$, such that each $\varkappa_n = E_n^+ - E_n^-$ $(n \in \mathbb{N})$. Consequently, from the gap lengths $(|\gamma_n|)_1^\infty$ one can uniquely recover the Riemann surface Λ as well as the points $E_n^- = E_n^+$ where $\varkappa_n = 0$. Furthermore, for any additional sequence $\widetilde{\lambda}_n^0 \in \gamma_n^c$ $(n \in \mathbb{N})$,

there is a unique potential $p \in \mathcal{H}$ such that each state λ_n^0 of the corresponding operator coincides with $\widetilde{\lambda}_n^0$ $(n \in \mathbb{N})$. The results of [K5] were extended in [K6] to periodic distributions $p = w'$, where $w \in \mathcal{H}$.

Now let

$$D(\lambda) = \det(I + q(H_0 - \lambda)^{-1}) \qquad (\lambda \in \mathbb{C}_+).$$

The function D is analytic in $\mathbb{C}_+$ and has a meromorphic extension to Λ. Each zero of D in Λ_1 is an eigenvalue of H and lies somewhere in the union of the physical gaps $\bigcup_{n\in\mathbb{N}_0} \gamma_n^{(1)}$. So far only certain particular results are known concerning the zeros on the non-physical sheet Λ_2. Note that the set of zeros of D on Λ_2 is symmetric with respect to the real line, since D is real on $\gamma_0^{(2)}$.

We now turn to the perturbed equation. Let $\Phi(x, z)$ be the solution of the initial-value problem

$$-\Phi'' + (p+q)\Phi = z^2\Phi \text{ on } [0,\infty), \qquad \Phi(0,z) = 0, \quad \Phi'(0,z) = 1 \quad (z \in \mathbb{C}). \tag{1.6}$$

Then our first result is as follows.

Theorem 1.1. *i) Let* $\Omega_\varepsilon := \{z \in \mathbb{C} \mid |z^2 - E_n^\pm| \geqslant n\varepsilon \ (n \in \mathbb{N})\}$ *for some* $\varepsilon > 0$. *Then the function* D *satisfies*

$$D(z^2) = 1 + \frac{\widehat{q}(z) - \widehat{q}(0)}{2iz} + \frac{O(e^{t(|\operatorname{Im} z| - \operatorname{Im} z)})}{z^2} \qquad \text{as} \quad |z| \to \infty, z \in \Omega_\varepsilon \tag{1.7}$$

where $\widehat{q}(z) = \int_0^t q(x) e^{2izx} dx$, *and*

$$\mathfrak{S}_{st}(H) \setminus \mathfrak{S}_{st}(H_0) = \{\lambda \in \Lambda \setminus \mathfrak{S}_{st}(H_0) \mid \ D(\lambda) = 0\} \subset \Lambda_2 \cup \bigcup_{n\in\mathbb{N}_0} \gamma_n^{(1)}, \tag{1.8}$$

$$\mathfrak{S}_{st}(H) \cap \mathfrak{S}_{st}(H_0) = \{z \in \mathfrak{S}_{st}(H_0) \mid \Phi(n_t, z) = 0\}. \tag{1.9}$$

ii) If $\lambda_n^0 \in \mathfrak{S}_{st}(H) \cap \mathfrak{S}_{st}(H_0)$, *then* $\lambda_n^0 \in \gamma_n^c \neq \emptyset$ $(n \in \mathbb{N})$ *and*

$$D(\lambda) \to D(\lambda_n^0) \neq 0 \quad \text{as} \quad \lambda \to \lambda_n^0. \tag{1.10}$$

iii) (Logarithmic Law) Each resonance $\lambda \in \Lambda_2$ *of* H *satisfies*

$$|\sqrt{\lambda} \sin \sqrt{\lambda}| \leqslant C_F e^{(2t+1)|\operatorname{Im}\sqrt{\lambda}|}, \qquad C_F = 3(\|p\|_1 + \|p+q\|_t) e^{2\|p+q\|_t + \|p\|_1}, \tag{1.11}$$

and there are no resonances in the domain $\mathscr{D}_{forb} = \{\lambda \in \Lambda_2 \setminus \cup \overline{\gamma}_n^{(2)} \mid 4C_F e^{2|\operatorname{Im}\sqrt{\lambda}|} < |\lambda|^{\frac{1}{2}}\}$. *Here* $\|f\|_s := \int_0^s |f(x)|\, dx$ $(s > 0)$.

Remarks. 1) Let $\lambda_n^0 \in \gamma_n^{(1)}$ be an eigenvalue of H_0 for some $n \geqslant 1$. If $\Phi(n_t, \mu_n) = 0$, then μ_n^2 is an eigenvalue of the Dirichlet boundary value problem $-y'' + (p+q)y = \mu_n^2 y, y(0) = y(n_t) = 0$. Then by (1.9), λ_n^0 is a bound state of H and (1.10) yields $D(\lambda_n^0) \neq 0$. Thus λ_n^0 **is a pole of a resolvent, but** λ_n^0 **is neither a zero of** D **nor a pole** of the S-matrix for H, H_0 given by

$$\mathcal{S}_M(z) = \frac{\overline{D(\lambda)}}{D(\lambda)}, \quad \lambda \in \sigma_{ac}(H_0). \tag{1.12}$$

2) If $D(\lambda) = 0$ for some $\lambda = E_n^\pm \neq \mu_n^2$, $n \in \mathbb{N}_0$, then by (1.8), λ is a virtual state.

3) If $\mu_n^2 = E_n^\pm$ for some $n \in \mathbb{N}$, then by (1.9), μ_n^2 is a virtual state if and only if $\Phi(n_t, \mu_n) = 0$.



FIGURE 2. The eigenvalues on the physical gaps $\gamma_n^1 \subset \Lambda_1$, anti-bound states on the non-physical gaps $\gamma_n^2 \subset \Lambda_2$, resonances on the non-physical sheet Λ_2 and the forbidden domain $\mathscr{D}_{forb}$

4) If $p = 0$, then it is well known that each zero of D is a state ([K1], [S2]). Moreover, each resonance lies below a logarithmic curve depending only on q ([K1], [Z]). The forbidden domain $\mathscr{D}_{forb} \cap \mathbb{C}_-$ is similar to the one in the case $p = 0$, see [K1].

Let $\#(H, r, A)$ be the total number, counted according to multiplicity, of states of H of modulus $\leqslant r$ in the set $A \subseteq \Lambda$.

The Fourier coefficients $\widehat{p}_{sn}, \widehat{q}_{cn}$ and the Fourier transform $\widehat{q}$ are defined by

$$\widehat{q}_0 = \int_0^t q(x)dx, \quad \widehat{p}_{sn} = \int_0^1 p(x)\sin 2\pi n x dx, \quad \widehat{q}(z) = \int_0^t q(x)e^{2izx}dx, \quad \widehat{q}_{cn} = \operatorname{Re}\widehat{q}(\pi n). \quad (1.13)$$

Theorem 1.2. *i) H has an odd number $\geqslant 1$ of states on each set $\gamma_n^c \neq \emptyset, n \geqslant 1$, where γ_n^c is the union of the physical gap $\overline{\gamma}_n^{(1)} \subset \Lambda_1$ and the non-physical gap $\overline{\gamma}_n^{(2)} \subset \Lambda_2$, and H has exactly one simple state $\lambda_n \in \gamma_n^c$ for all $n \geqslant 1 + 4C_F e^{t\frac{\pi}{2}}$ with asymptotics*

$$\sqrt{\lambda_n} = \mu_n - \frac{(\widehat{q}_0 - \widehat{q}_{cn})\widehat{p}_{sn}}{2(\pi n)^2} + O(\frac{1}{n^3}) \qquad \text{as} \quad n \to \infty. \quad (1.14)$$

Moreover, the following asymptotics hold true as $r \to \infty$:

$$\#(H, r, \Lambda_2 \setminus \cup\gamma_n^{(2)}) = r\frac{2t}{\pi} + o(r), \quad (1.15)$$

$$\#(H, r, \gamma_\Lambda) = \#(H_0, r, \gamma_\Lambda) + 2N_q \qquad for\ some\ integer\ N_q \geqslant 0, \quad r \notin \cup\overline{\gamma}_n, \quad (1.16)$$

where $\gamma_\Lambda = \bigcup_{n\geqslant 0} \overline{\gamma}_n^{(1)} \cup \overline{\gamma}_n^{(2)}$ is the union of all gaps on Λ.
ii) Let λ be an eigenvalue of H and let $\lambda^{(2)} \in \Lambda_2$ be the same number but on the second sheet Λ_2. Then $\lambda^{(2)}$ is not an anti-bound state.

iii) Let $\lambda_1, \lambda_2 \in \gamma_n^{(1)}, \lambda_1 < \lambda_2$*, be eigenvalues of* H *for some* $n \geqslant 0$ *and assume that there are no other eigenvalues on the interval* $\Omega = (\lambda_1, \lambda_2) \subset \gamma_n^{(1)}$*. Let* $\Omega^{(2)} \subset \gamma_n^{(2)} \subset \Lambda_2$ *be the same interval but on the second sheet. Then there exists an odd number* $\geqslant 1$ *of antibound states on* $\Omega^{(2)}$*.*

Remarks. 1) Results (iii) with $p = 0$ were obtained independently in [K1], [S2].

2) The first term in the asymptotics (1.15) is independent of the periodic potential p. The asymptotics (1.15) for the case $p = 0$ was obtained by Zworski [Z].

3) The main difference between the distribution of the resonances in the cases $p \neq$ const and $p =$ const concerns the bound states and antibound states in high energy gaps, see (1.14).

4) In the proof of (1.15) we use a Paley-Wiener type theorem from [Fr], the Levinson Theorem (see Sect. 4) and an analysis of the function D near λ_n^0.

5) For even potentials $p \in L^2_{even}(0, 1) = \{p \in L^2(0, 1) \mid p(x) = p(1 - x), x \in (0, 1)\}$, all coefficients p_{sn} vanish and the asymptotics (1.14) are not sharp. This case is analyzed further in Theorem 1.4.

We now turn to the question of stability of the real states λ_n. As before, let $\lambda_n^0 \in \gamma_n^c$ be a state of H_0.

Theorem 1.3. *Let* $b_n = q_0 - \widehat{q}_{cn}$ $(n \in \mathbb{N})$*. Assume that* $|p_{sn}| > n^{-\alpha}$ *and* $|b_n| > n^{-(1-\alpha)}$ *for some* $\alpha \in (0, 1)$ *and for all* $n \in \mathcal{N}_0$*, where* $\mathcal{N}_0 \subset \mathbb{N}$ *is some infinite subset such that* $|\gamma_n| > 0$ $(n \in \mathcal{N}_0)$*. Assume* $b_n > 0$ *(or* $b_n < 0$*) for all* $n \in \mathcal{N}_0$*.*

Then each real state $\lambda_n \in \gamma_n^c$ *of the perturbed operator* H *has, for sufficiently large* $n \in \mathcal{N}_0$*, the following property.*
If λ_n^0 *is an eigenvalue of* H_0*, then* λ_n *is an eigenvalue of* H *and* $\lambda_n^0 < \lambda_n$ *(or* $\lambda_n^0 > \lambda_n$*).*
If λ_n^0 *is an antibound state of* H_0*, then* λ_n *is an antibound state of* H *and* $\lambda_n^0 > \lambda_n$ *(or* $\lambda_n^0 < \lambda_n$*).*

Remarks. 1) Let $q > 0$. It is well known that the eigenvalues of $H_0 + \tau q$ are increasing with the coupling constant $\tau > 0$. Roughly speaking, in the case considered in Theorem 1.3 the antibound states in the gap move in the opposite direction.

2) Numerical observations suggest the following scenario as the coupling constant of the perturbation changes. Consider the operator $H_\tau = H_0 + \tau q$, where $\tau \in \mathbb{R}$ is the coupling constant. For $\tau = 0$, H_0 only has states λ_n^0, $n \in \mathbb{N}$ (eigenvalues, antibound states and virtual states). Take for instance the first gap $\gamma_1^c \neq \emptyset$ and assume that $\lambda_1 = \lambda_1^0$ is an antibound state. As τ increases, the state λ_1 moves, and initially there are no other states on γ_1^c. As τ increases further, λ_1 eventually emerges onto the physical gap $\gamma_1^{(1)}$ and becomes an eigenvalue; at first there are no additional eigenvalues, but a pair of complex resonances ($\lambda \in \mathbb{C}_+ \subset \Lambda_2$ and $\overline{\lambda} \in \mathbb{C}_- \subset \Lambda_2$) reach the non-physical gap $\gamma_1^{(2)}$ and there turn into antibound states. With further increase of τ, these antibound states will become virtual states and then bound states. Thus, as τ runs through $\mathbb{R}_+$, states not only move, but are also subject to the following transmutations: resonances turn into antibound states, these via virtual states into bound states, these via virtual states into antibound states etc.

We now consider the perturbations of virtual states. Note that $p \in L^2_{even}(0, 1)$ if and only if $\mu_n^2 \in \{E_n^-, E_n^+\}$ for all $n \in \mathbb{N}$ ([GT], [KK1]).

Theorem 1.4. *Let $p \in L^1(0,1)$ and assume there are unperturbed states $\lambda_n^0 \in \{E_n^-, E_n^+\}$ for all $n \in \mathcal{N}_0$, where $\mathcal{N}_0 \subset \mathbb{N}$ is some infinite subset such that $|\gamma_n| > 0$ $(n \in \mathcal{N}_0)$. Then the following asymptotics for states of H hold true:*

$$\sqrt{\lambda_n} = \mu_n + s_n|\gamma_n|\frac{(q_0 - \widehat{q}_{cn} + O(\frac{1}{n}))^2}{(2\pi n)^2}, \qquad s_n = \begin{cases} + & \text{if } \mu_n^2 = E_n^- \\ - & \text{if } \mu_n^2 = E_n^+ \end{cases}, \quad n \in \mathcal{N}_0 \tag{1.17}$$

as $n \to \infty$. Moreover, if $\alpha \in (\frac{1}{2}, 1)$, then for sufficiently large $n \in \mathcal{N}_0$, the following holds:
if $\lambda_n^0 = E_n^-$, $q_0 - \widehat{q}_{cn} > n^{-\alpha}$ or $\lambda_n^0 = E_n^+$, $q_0 - \widehat{q}_{cn} < -n^{-\alpha}$, then λ_n is an eigenvalue,
if $\lambda_n^0 = E_n^-$, $q_0 - \widehat{q}_{cn} < -n^{-\alpha}$ or $\lambda_n^0 = E_n^+$, $q_0 - \widehat{q}_{cn} > n^{-\alpha}$, then λ_n is an antibound state.

Remark. 1) Roughly speaking, (1.17) gives the asymptotics for even potentials $p \in L^1(0,1)$.

2) Consider the operator $-y''+(p+q+u)y$, $y(0)=0$, with an additional potential perturbation $u \in L^2(\mathbb{R}_+)$ which is compactly supported in $(0,t)$ and satisfies $|\widehat{u}_n| = o(n^{-1})$ as $n \to \infty$. Then the operator $H+u$ has the same number of bound states as H in each gap $\gamma_n \neq \emptyset$ for n large enough.

We have the following result on the inverse problem for our operator H.

Theorem 1.5. *i) Let $q \in \mathcal{Q}_t$ satisfy $|q_0 - \widehat{q}_{cn}| > n^{-\alpha}$ for sufficiently large n, with some $\alpha \in (\frac{1}{2}, 1)$. Then for any sequences $(\sigma_n)_1^\infty, \sigma_n \in \{0,1\}$, and $(\varkappa_n)_1^\infty \in \ell^2, \varkappa_n \geqslant 0$, there exists a potential $p \in L^2(0,1)$ such that the corresponding gap lengths $|\gamma_n|$ satisfy $|\gamma_n| = \varkappa_n, n \geqslant 1$, and H has exactly σ_n eigenvalues and $1-\sigma_n$ antibound states in each gap $\gamma_n \neq \emptyset$ for sufficiently large n.*
ii) Let $p \in L^1(0,1)$ and assume there are unperturbed states $\lambda_n^0 \in \{E_n^-, E_n^+\}$ for all $n \in \mathcal{N}_0$, where $\mathcal{N}_0 \subset \mathbb{N}$ is some infinite subset such that $|\gamma_n| > 0$ $(n \in \mathcal{N}_0)$. Then for any sequence $(\sigma_n)_1^\infty, \sigma_n \in \{0,1\}$, there exists a potential $q \in \mathcal{Q}_t$ such that H has exactly σ_n eigenvalues and $1-\sigma_n$ antibound states in each gap $\gamma_n \neq \emptyset$ for sufficiently large $n \in \mathcal{N}_0$.

Let $\#_{bs}(H,\Omega)$ $(\#_{abs}(H,\Omega))$ be the total number, counted according to multiplicity, of bound (antibound) states of H on the segment $\Omega \subset \overline{\gamma}_n^{(1)} \subset \Lambda_1$ $(\Omega \subset \overline{\gamma}_n^{(2)} \subset \Lambda_2)$ for some $n \in \mathbb{N}_0$.

The integrated density of states ϱ can be characterized as a continuous, real-valued function on $\mathbb{R}$ with the properties

$$\varrho([E_n^-, E_n^+]) = n, \qquad \varrho(\mathfrak{S}_{n+1}) = [n, n+1], \qquad \cos\pi\varrho(\lambda) = \Delta(\sqrt{\lambda}) \ (\lambda \in \mathfrak{S}_{n+1}) \tag{1.18}$$

for all $n \in \mathbb{N}_0$. The function ϱ is strictly increasing on each spectral band $\mathfrak{S}_n$ and constant on the closure of each gap $[E_n^-, E_n^+]$. It is closely related to the quasimomentum defined in Section 2 via $\varrho(\lambda) = \frac{1}{\pi}\operatorname{Re} k(\sqrt{\lambda + i0})$ $(\lambda \in \mathbb{R})$. Theorem 1.2 yields the following corollary.

Theorem 1.6. *Let $H_\tau = H_0 + q_\tau$ where $q_\tau = q(\frac{\cdot}{\tau}), \tau \geqslant 1$. Let $\Omega = [E_1, E_2] \subset \overline{\gamma}_n^{(1)} \neq \emptyset$ be some interval on the physical sheet Λ_1 for some $n \geqslant 0$ and let $\Omega^{(2)} \subset \overline{\gamma}_n^{(2)}$ be the corresponding interval on the non-physical sheet Λ_2. Then*

$$\begin{aligned} \#_{abs}(H_\tau, \Omega^{(2)}) &\geqslant 1 + \#_{bs}(H_\tau, \Omega) \\ &= \tau \int_0^\infty \Big(\varrho(E_2 - q(x)) - \varrho(E_1 - q(x)) \Big) dx + o(\tau) \quad \text{as} \quad \tau \to \infty. \end{aligned} \tag{1.19}$$

Remark. 1) The proof of (1.19) will be based on Sobolev's idea [So] for obtaining asymptotics of $\#_{bs}(H_\tau, \Omega)$. He considered the case $H_\tau = H_0 + \tau V$, where $V(x) = \frac{c+o(1)}{x^\alpha}$ as $x \to \infty$, for some $c \neq 0, \alpha > 0$. This result is not directly applicable to our case of a compactly supported perturbation. We therefore follow the reinterpretation of the coupling constant as a scaling parameter, as previously used in the study of perturbations of the periodic Dirac operator [Sc], which allows compactly supported perturbations.

2) Symmetry of bound states and antibound states for $p = 0$ in the semiclassical limit was studied in [BZ], [DG].

A large number of papers are devoted to resonances for the Schrödinger operator with $p = 0$, see [Fr], [H], [K1], [K2], [S2], [Z] and references therein. Although resonances have been studied in many settings, there are relatively few cases in which the asymptotics of the resonance counting function are known, mainly in the one-dimensional setting ([Fr], [K1], [K2], [S2], and [Z]). Zworski [Z] obtained the first results about the distribution of resonances for the Schrödinger operator with compactly supported potentials on the real line. One of the present authors obtained the characterization, including uniqueness and recovery, of the S-matrix for the Schrödinger operator with a compactly supported potential on the real line [K2] and on the half-line [K1], see also [Z1], [BKW] concerning the uniqueness.

The following stability results for the Schrödinger operator on the half line can be found in [K3]; here the real Hilbert spaces ℓ^2_ε are given by

$$\ell^2_\varepsilon = \Big\{ f = (f_n)_1^\infty \mid \ f_n \in \mathbb{R}, \quad \|f\|^2_\varepsilon = \sum_{n \geqslant 1} (2\pi n)^{2\varepsilon} f_n^2 < \infty \Big\} \ \ (\varepsilon \geqslant 0), \qquad \ell^2_0 = \ell^2.$$

(i) If $\varkappa = (\varkappa)_1^\infty$ is a sequence of poles (eigenvalues and resonances) of the S-matrix for some real compactly supported potential q and $\widetilde{\varkappa} - \varkappa \in \ell^2_\varepsilon$ for some $\varepsilon > 1$, then $\widetilde{\varkappa}$ is a sequence of poles of the S-matrix for some uniquely determined real-valued compactly supported potential $\widetilde{q}$.
(ii) The measure associated with the poles of the S-matrix is the Carleson measure, and the sum $\sum (1 + |\varkappa_n|)^{-\alpha}, \alpha > 1$, can be estimated in terms of the L^1-norm of the potential q.

Brown and Weikard [BW] considered the Schrödinger operator $-y'' + (p_A + q)y$ on the half-line, where p_A is an algebro-geometric potential and q is a compactly supported perturbation. They proved that the zeros of the Jost function determine q uniquely.

Christiansen [Ch] considered resonances associated with the Schrödinger operator $-y'' + (p_S + q)y$ on the real line, where p_S is a step potential. She determined the asymptotics of the resonance-counting function and showed that the resonances determine q uniquely.

The recent paper [K4] gives the following results about the operator $\mathscr{H} = \mathscr{H}_0 + q$, $\mathscr{H}_0 = -\frac{d^2}{dx^2} + p$ on the real line, where p is periodic and q is compactly supported: 1) asymptotics of the resonance-counting function are determined, 2) a forbidden domain for the resonances is specified, 3) the asymptotics of eigenvalues and antibound states are determined, 4) for any sequence $(\sigma)_1^\infty, \sigma_n \in \{0, 2\}$, there exists a compactly supported potential q such that $\mathscr{H}$ has σ_n bound states and $2 - \sigma_n$ antibound states in each gap $\gamma_n \neq \emptyset$ for n large enough, 5) for any q (with $q_0 = 0$) and for any sequences $(\sigma_n)_1^\infty, \sigma_n \in \{0, 2\}$, and $(\varkappa_n)_1^\infty \in \ell^2, \varkappa_n \geqslant 0$, there exists a potential $p \in L^2(0,1)$ such that gap γ_n has length $|\gamma_n| = \varkappa_n$ ($n \in \mathbb{N}$) and $\mathscr{H}$ has exactly σ_n eigenvalues and $2 - \sigma_n$ antibound states in each gap $\gamma_n \neq \emptyset$ for sufficiently large n.



FIGURE 3. The cut domain $\mathscr{Z} = \mathbb{C} \setminus \cup \overline{g}_n$ and the slits $g_n = (e_n^-, e_n^+)$ in the z-plane.

Thus, comparing the results for $\mathscr{H}$ on $\mathbb{R}$ and H on $\mathbb{R}_+$, we find that their properties are close for even potentials $p \in L^2_{even}(0,1)$, since in this case the unperturbed operator H_0 has only virtual states; however, if p is not even, then the unperturbed operator H_0 has eigenvalues, virtual states and antibound states in general, but the operator $\mathscr{H}_0$ has exactly two virtual states in each non-empty gap. This leads to the different properties of the perturbed operators $H, \mathscr{H}$, and, roughly speaking, the case of H is more complicated since the set of states of the unperturbed operator H_0 is already more complicated.

For related results in the multidimensional case, see [G], [D] and references therein.

The structure of the present paper is as follows. In Section 2 we define the Riemann surface associated with the momentum variable $z = \sqrt{\lambda}, \lambda \in \Lambda$, and describe preliminary results about fundamental solutions. In Section 3 we study the states of H. In Section 4 we prove the main Theorems 1.1-1.6. The proofs use properties of the quasimomentum, a priori estimates from [KK], [MO], and results from the inverse theory for the Hill operator [K5]. The analysis of the entire function $F(z) = \varphi(1,z) D(z) \overline{D}(z), z^2 \in \sigma_{ac}(H)$, plays an important role, since its zeros are, as we show, closely related to the states. Thus the spectral analysis of H is reduced to a problem of entire function theory.

2. PRELIMINARIES

In the following, we shall treat the momentum $z = \sqrt{\lambda}$ (as opposed to the energy λ) as the principal spectral variable. Note that the functions $\varphi(x,z), \vartheta(x,z)$ are entire in $z \in \mathbb{C}$ (cf. [E1] Thm 1.7.2), and so are the functions $\widetilde{\vartheta}, \widetilde{\varphi}$ defined in (2.16) below. The other functions considered in the present paper are combinations of $\varphi(x,z), \vartheta(x,z), \widetilde{\varphi}(x,z), \widetilde{\vartheta}(x,z)$ and $m_\pm = \frac{\beta \pm i \sin k}{\varphi(1,\cdot)}$. Thus we are led to consider the Riemann surface $\mathcal{M}$ corresponding to the analytic continuation of the function $\sin k$ (cf. [K4]). Take the cut domain (see Fig.3)

$$\mathscr{Z} = \mathbb{C} \setminus \cup \overline{g}_n, \qquad \text{where} \quad g_n = (e_n^-, e_n^+) = -g_{-n}, \quad e_n^\pm = \sqrt{E_n^\pm} > 0, \quad n \geqslant 1, \tag{2.1}$$

and note that $\Delta(e_n^\pm) = (-1)^n$. If $\lambda \in \gamma_n, n \geqslant 1$, then $z \in g_{\pm n}$, and if $\lambda \in \gamma_0 = (-\infty, 0)$, then $z \in g_0^\pm = i\mathbb{R}_\pm$. Slitting the n-th momentum gap g_n (if non-empty), we obtain a cut g_n^c with an upper rim g_n^+ and lower rim g_n^-. Below we will identify this cut g_n^c and the union of the upper rim (gap) $\overline{g}_n^+$ and the lower rim (gap) $\overline{g}_n^{\,-}$, i.e.,

$$g_n^c = \overline{g}_n^+ \cup \overline{g}_n^-, \text{ where } g_n^\pm = g_n \pm i0; \quad \text{and } z \in g_n \Rightarrow z \pm i0 \in g_n^\pm. \tag{2.2}$$

Now the function $\sin k(z) = \sqrt{1 - \Delta^2(z)}, z \in \mathscr{Z}$, which is analytic because $\varphi(x,z), \vartheta'(x,z)$ are, has an analytic continuation through the cuts $\cup g_n \subset \mathbb{R}$, since $\sqrt{1 - \Delta^2(z)} \in i\mathbb{R}$ for all



FIGURE 4. The domain $\mathcal{K}=\mathbb{C}\setminus\cup\Gamma_n$ with $\Gamma_n=(\pi n-ih_n,\pi n+ih_n)$

$z\in\cup g_n$. Thus we obtain the analytic function $\sqrt{1-\Delta^2(z)}$ on the Riemann surface $\mathcal{M}$, and (by applying the square root function) the analytic function $\sqrt{1-\Delta^2(\sqrt{\lambda})}$ on the Riemann surface Λ.

The Riemann surface $\mathcal{M}$ can be obtained from the cut domain $\mathcal{Z}=\mathbb{C}\setminus\cup\overline{g}_n$ as follows. We identify the upper rim g_n^+ of the cut g_n^c with the upper rim g_{-n}^+ of the slit g_{-n}^c and, correspondingly, the lower rim g_n^- of the cut g_n^c with the lower rim g_{-n}^- of the cut g_{-n}^c for all non-empty gaps. The mapping $z=\sqrt{\lambda}$ gives a bijection from Λ onto $\mathcal{M}$. The gap $\gamma_n^{(1)}\subset\Lambda_1$ is mapped onto $g_n^+\subset\mathcal{M}$ and the gap $\gamma_n^{(2)}\subset\Lambda_2$ is mapped onto $g_n^-\subset\mathcal{M}$. The upper rim g_n^+ corresponds to a physical gap, the lower rim g_n^- to a non-physical gap. Moreover, the union of $\mathcal{M}\cap\mathbb{C}_+=\mathcal{Z}\cap\mathbb{C}_+$ with all physical gaps g_n^+ forms the so-called physical sheet $\mathcal{M}_1$, while the union of $\mathcal{M}\cap\mathbb{C}_-=\mathcal{Z}\cap\mathbb{C}_-$ with all non-physical gaps g_n^- forms the so-called non-physical sheet $\mathcal{M}_2$. The two sheets are joined together on the set $\mathbb{R}\setminus\cup g_n$ (which is the spectrum of the periodic operator H_0).

We introduce the *quasimomentum* k for H_0 as $k(z)=\arccos\Delta(z)$ $(z\in\mathcal{Z})$. The function k is analytic in $\mathcal{Z}$ and satisfies

$$(i)\quad k(z)=z+O(1/z)\quad as\quad |z|\to\infty,\qquad (ii)\quad \operatorname{Re}k(z\pm i0)|_{[e_n^-,e_n^+]}=\pi n\quad (n\in\mathbb{Z}),\tag{2.3}$$

as well as $\pm\operatorname{Im}k(z)>0$ for any $z\in\mathbb{C}_\pm$ ([MO], [KK]). The function sin of the quasimomentum k is also analytic on $\mathcal{M}$ and satisfies $\sin k(z)=(1-\Delta^2(z))^{\frac12}$ $(z\in\mathcal{M})$. Moreover, k is a conformal mapping from $\mathcal{Z}$ onto the quasimomentum domain $\mathcal{K}=\mathbb{C}\setminus\cup\Gamma_n$, see Figs. 3 and 4. Here $\Gamma_n=(\pi n-ih_n,\pi n+ih_n)$ is a vertical slit of height $2h_n\geqslant 0, h_0=0$. The height h_n is determined by the equation $\cosh h_n=(-1)^n\Delta(e_n)\geqslant 1$, where $e_n\in[e_n^-,e_n^+]$ is such that $\Delta'(e_n)=0$. The function k maps the slit g_n^c onto the slit Γ_n, and $k(-z)=-k(z)$ for all $z\in\mathcal{Z}$.

In order to describe the spectral properties of the operator H_0, we start from the properties of the canonical fundamental system ϑ,φ of the equation $-y''+py=z^2y$. They satisfy the integral equations

$$\begin{aligned}\vartheta(x,z)&=\cos zx+\int_0^x\frac{\sin z(x-s)}{z}p(s)\vartheta(s,z)ds,\\ \varphi(x,z)&=\frac{\sin zx}{z}+\int_0^x\frac{\sin z(x-s)}{z}p(s)\varphi(s,z)ds.\end{aligned}\tag{2.4}$$

For each fixed $x \in \mathbb{R}$, the functions $\vartheta(x,\cdot)$, $\varphi(x,\cdot)$ are entire in $\mathbb{C}$ and satisfy the estimates

$$\max\left\{|z|_1|\varphi(x,z)|,\ |\varphi'(x,z)|, |\vartheta(x,z)|, \frac{1}{|z|_1}|\vartheta'(x,z)|\right\} \leqslant X = e^{|\operatorname{Im} z|x+\|p\|_x},$$
$$\left|\varphi(x,z) - \frac{\sin zx}{z}\right| \leqslant \frac{X}{|z|^2}\|p\|_x, \quad |\vartheta(x,z) - \cos zx| \leqslant \frac{X}{|z|}\|p\|_x, \quad (2.5)$$

where $|z|_1 = \max\{1,|z|\}$, $\|p\|_x = \int_0^x |p(s)|ds$ and $(x,z) \in \mathbb{R}\times\mathbb{C}$, see page 13 in [PT]. Substituting these estimates into (2.4) we obtain the standard asymptotics

$$\beta(z) = \frac{\varphi'(1,z) - \vartheta(1,z)}{2} = \int_0^1 \frac{\sin z(2x-1)}{z} p(x)dx + \frac{O(e^{|\operatorname{Im} z|})}{z^2} \qquad \text{as} \qquad |z| \to \infty. \quad (2.6)$$

Moreover, if $z = \pi n + O(1/n)$, then we obtain

$$\beta(z) = (-1)^n \frac{p_{sn}}{2\pi n} + O(1/n^2), \quad p_{sn} = \int_0^1 p(x)\sin 2\pi n x dx. \quad (2.7)$$

The Floquet solutions $\psi^{\pm}(x,z), z \in \mathcal{Z}$, of the equation $-y'' + py = z^2 y$ are given by

$$\psi^{\pm}(x,z) = \vartheta(x,z) + m_{\pm}(z)\varphi(x,z), \qquad m_{\pm} = \frac{\beta \pm i \sin k}{\varphi(1,\cdot)}, \quad (2.8)$$

where $\varphi(1,z)\psi^+(\cdot,z) \in L^2(\mathbb{R}_+)$ for all $z \in \mathbb{C}_+ \cup \bigcup_{n\in\mathbb{Z}} g_n$. In the trivial case $p = 0$, we have $k = z$ and $\psi^{\pm}(x,z) = e^{\pm izx}$.

Denote by $\mathcal{D}_r(z_0) = \{|z - z_0| < r\}$ the disc of radius $r > 0$ centred at $z_0 \in \Lambda$. It is well known that if $g_n = \emptyset$ for some $n \in \mathbb{Z}$, then the functions $\sin k$ and $m_{\pm}$ *are analytic in the disc* $\mathcal{D}_\varepsilon(\mu_n) \subset \mathcal{Z}$ for some $\varepsilon > 0$, and the functions $\sin k$ and $\varphi(1,\cdot)$ have a simple zero at μ_n [F1]. Moreover, $m_{\pm}$ satisfies

$$m_{\pm}(\mu_n) = \frac{\beta'(\mu_n) \pm i(-1)^n k'(\mu_n)}{\partial_z \varphi(1,\mu_n)}, \quad \operatorname{Im} m_{\pm}(\mu_n) \neq 0. \quad (2.9)$$

Furthermore, $\operatorname{Im} m_+(z) > 0$ for all $z \in (z_{n-1}^+, z_n^-), n \in \mathbb{N}$, and the following asymptotics hold,

$$m_{\pm}(z) = \pm iz + O(1) \quad as \quad |z| \to \infty, \quad z \in \mathcal{Z}_\varepsilon,\ \varepsilon > 0, \quad (2.10)$$

where $\mathcal{Z}_\varepsilon = \{z \in \mathcal{Z} \mid \operatorname{dist}\{z, g_n\} > \varepsilon\ (n \in \mathbb{Z}, g_n \neq \emptyset)\}$. The function $\sin k$ and each function $\varphi(1,\cdot)\psi^{\pm}(x,\cdot)$, $x \in \mathbb{R}$, are analytic on the Riemann surface $\mathcal{M}$. From (2.8), the Floquet solutions $\psi^{\pm}(x,z)$ $((x,z) \in \mathbb{R}\times\mathcal{M})$ satisfy [T]

$$\psi^{\pm}(0,z) = 1, \quad \psi^{\pm}(0,z)' = m_{\pm}(z), \quad \psi^{\pm}(1,z) = e^{\pm ik(z)}, \quad \psi^{\pm}(1,z)' = e^{\pm ik(z)} m_{\pm}(z), \quad (2.11)$$

$$\psi^{\pm}(x,z) = e^{\pm ik(z)x}(1 + O(1/z)) \quad (2.12)$$

as $|z| \to \infty, z \in \mathcal{Z}_\varepsilon$, uniformly in $x \in \mathbb{R}$. Below we shall require the simple identities

$$\beta^2 + 1 - \Delta^2 = 1 - \varphi'(1,\cdot)\vartheta(1,\cdot) = -\varphi(1,\cdot)\vartheta'(1,\cdot). \quad (2.13)$$

Now consider the modified Jost solutions $\Psi^{\pm}(x,z)$ of the equation

$$-\Psi^{\pm\prime\prime} + (p+q)\Psi^{\pm} = z^2\Psi^{\pm} \text{ on } [0,\infty) \quad \text{with } \Psi^{\pm}(x,z) = \psi^{\pm}(x,z),\ (x \geqslant t, z \in \mathcal{Z}\setminus\{0\}). \quad (2.14)$$

Each function $\varphi(1,\cdot)\Psi^{\pm}(x,\cdot)$, $x \geqslant 0$, is analytic in $\mathcal{M}$, since each $\varphi(1,\cdot)\psi^{\pm}(x,\cdot)$, $x \geqslant 0$, is analytic in $\mathcal{M}$. We define the *modified Jost function* $\Psi_0^{\pm} = \Psi^{\pm}(0,z)$, which is meromorphic in $\mathcal{M}$ and has branch points $e_n^{\pm}$ $(n \in \mathbb{Z}, g_n \neq \emptyset)$.

Let $\varphi(x,z,\tau)$ $((z,\tau)\in\mathbb{C}\times\mathbb{R})$ be the solutions of the initial-value problem

$$-\varphi''+p(x+\tau)\varphi=z^2\varphi,\qquad \varphi(0,z,\tau)=0,\quad \varphi'(0,z,\tau)=1. \tag{2.15}$$

The solutions $\widetilde{\vartheta},\widetilde{\varphi}$ of the perturbed equation $-y''+(p+q)y=z^2y, z\in\mathbb{C}$, determined by

$$\widetilde{\vartheta}(x,z)=\vartheta(x,z),\ \widetilde{\varphi}(x,z)=\varphi(x,z)\ (\quad x\geqslant t),$$

satisfy the integral equations

$$\begin{aligned}\widetilde{\vartheta}(x,z)&=\vartheta(x,z)-\int_x^t\varphi(x-s,z,s)q(s)\widetilde{\vartheta}(s,z)\,ds,\\ \widetilde{\varphi}(x,z)&=\varphi(x,z)-\int_x^t\varphi(x-s,z,s)q(s)\widetilde{\varphi}(s,z)\,ds.\end{aligned} \tag{2.16}$$

For each $x\geqslant 0$, $\widetilde{\vartheta}(x,z)$, $\widetilde{\varphi}(x,z)$ are entire functions of z and take real values if $z^2\in\mathbb{R}$. The identities (2.16) and (2.14) give

$$\Psi^{\pm}(x,z)=\widetilde{\vartheta}(x,z)+m_{\pm}(z)\widetilde{\varphi}(x,z),\ \ ((x,z)\in\mathbb{R}_+\times\mathscr{Z}). \tag{2.17}$$

We see that all singularities of $\Psi^{\pm}(x,z)$ coincide with some singularity of $m_{\pm}(z)$ and hence do not depend on $x>0$. The kernel of the resolvent $R=(H-z^2)^{-1}, z\in\mathbb{C}_+$, then has the form

$$R(x,x',z)=\frac{\Phi(x,z)\Psi^+(x',z)}{\Psi_0^+(z)}\ (x<x'),\ \text{and}\ R(x,x',z)=R(x',x,z)\ (x>x'). \tag{2.18}$$

Here $\Phi(x,z)$ is the solution of the perturbed equation defined in (1.6). Each function $R(x,x',z)$, $x,x'\in\mathbb{R}$, is meromorphic in $\mathcal{M}$. Note that if $z_0\in g_n^{\pm}\setminus\{\mu_n\pm i0\}$ for some n and $\Psi_0^+(z_0)\neq 0$, then the resolvent of H is analytic at z_0. The function Ψ_0^+ has a finite number of simple zeros on each $g_n^+, n\neq 0$, and on $i\mathbb{R}_+$, but no zeros on $\mathbb{C}_+\setminus i\mathbb{R}_+$, and the square of each zero is an eigenvalue. Note that the poles of

$$\mathscr{R}(x,z)=\frac{\Psi^+(x,z)}{\Psi^+(0,z)}=\frac{\widetilde{\vartheta}(x,z)+m_+(z)\widetilde{\vartheta}(x,z)}{\widetilde{\vartheta}(0,z)+m_+(z)\widetilde{\vartheta}(0,z)}$$

are either poles of m_+ or zeros of Ψ_0^+ and therefore are independent of x. A pole of $\mathscr{R}(x,\cdot)=\Psi^+(x,\cdot)/\Psi_0^+$ on g_n^+ is called a *bound state* by slight abuse of terminology (since it is a momentum rather than an energy value). Moreover, $\Psi_0^+(z)$ has an infinite number of zeros in $\mathbb{C}_-$, see (1.16). In terms of the Riemann surface $\mathcal{M}$, the previous definition of resonances and states, which referred to the Riemann surface Λ, takes the following form.

Definition. 1) A point $\zeta\in\overline{\mathbb{C}}_-\cap\mathcal{M},\zeta\neq 0$, is a *resonance* if the function $\mathscr{R}(x,z)$ has a pole at ζ for almost all $x>0$.
2) A point $\zeta=e_n^{\pm}, n\neq 0$, (or $\zeta=0$) is a *virtual state* if the function $z\mapsto\mathscr{R}(x,\zeta+z^2)$ (or $\mathscr{R}(x,z)$) has a pole at 0 for almost all $x>0$.
3) A point $\zeta\in\mathcal{M}$ is a *state* if it is either a bound state or a resonance or a virtual state. If $\zeta\in g_n^-, n\neq 0$, or $\zeta\in g_0^-=i\mathbb{R}_-$, then we call ζ an *antibound state.*

Let $\sigma_{bs}(H)$ (or $\sigma_{rs}(H)$ or $\sigma_{vs}(H)$) be the set of all bound states (or resonances or virtual states) of H in $\mathcal{M}$ (i.e., in terms of the momentum variable), and set $\sigma_{st}(H)=\sigma_{vs}(H)\cup\sigma_{rs}(H)\cup\sigma_{bs}(H)$.

The kernel of the resolvent $R_0(z) = (H_0 - z^2)^{-1}, z \in \mathbb{C}_+$, has the form

$$R_0(x,x',z) = \varphi(x,z)\psi^+(x',z)\ (x < x'), \quad \text{and} \quad R_0(x,x',z) = R_0(x',x,z)\ (x > x'). \tag{2.19}$$

Consider the states $z_n^0 = \sqrt{\lambda_n^0} \in g_n^c$ of H_0. Due to (2.8), the function $\Psi_0^+ = \psi^+(0,\cdot) = 1$ and $\mathscr{R}(x,\cdot) = \vartheta(x,\cdot) + m_+\varphi(x,\cdot)$. Since $\varphi(1,\cdot)m_+$ and $\sin k(z)$ are analytic in $\mathcal{M}$, the resolvent $R_0(z)$ has singularities only at $\mu_n \pm i0$, where $g_n \neq \emptyset$, and in order to describe the states of H_0 we need to study m_+ on g_n^c only. We need the following result (see [Zh3]), where $\mathscr{A}(z_0), z_0 \in \mathcal{M}$, denotes the set of functions analytic in some disc $\mathcal{D}_r(z_0), r > 0$.

Lemma 2.1. *All states of H_0 are of the form $\mu_n \pm i0 \in g_n^c, n \neq 0$, where $g_n \neq \emptyset$. Consider a non-empty momentum gap $g_n = (e_n^-, e_n^+)$ for some $n \geqslant 1$. Then*
i) $z_n^0 = \mu_n + i0 \in g_n^+$ is a bound state of H_0 if and only if one of the following conditions is true.

$$\begin{aligned}
&(1) \quad m_- \in \mathscr{A}(\mu_n + i0),\\
&(2) \quad \beta(\mu_n) = i \sin k(\mu_n + i0) = -(-1)^n \sinh h_{sn}, \quad k(\mu_n + i0) = \pi n + i h_{sn} \quad h_{sn} > 0,\\
&(3) \quad m_+(z_n^0 + z) = \frac{c_n}{z} + O(1) \quad \text{as } z \to 0,\ z \in \mathbb{C}_+,\ c_n = \frac{-2\sinh|h_{sn}|}{(-1)^n \partial_z \varphi(1,\mu_n)} < 0.
\end{aligned} \tag{2.20}$$

ii) $z_n^0 = \mu_n - i0 \in g_n^-$ is an antibound state of H_0 if and only if one of the following conditions is true.

$$\begin{aligned}
&(1) \qquad m_- \in \mathscr{A}(\mu_n - i0),\\
&(2) \qquad \beta(\mu_n) = i \sin k(\mu_n - i0) = -(-1)^n \sinh h_{sn}, \quad k(\mu_n - i0) = \pi n + i h_{sn}, \quad h_{sn} < 0,\\
&(3) \qquad m_+(z_n^0 + z) = \frac{-c_n}{z} + O(1) \quad \text{as } z \to 0, \quad z \in \mathbb{C}_-.
\end{aligned} \tag{2.21}$$

iii) $z_n^0 = \mu_n$ is a virtual state of H_0 if and only if one of the following conditions is true.

$$\begin{aligned}
&(1) \qquad z_n^0 = \mu_n = e_n^- \qquad \text{or} \qquad z_n^0 = \mu_n = e_n^+,\\
&(2) \qquad m_+(z_n^0 + z) = \frac{c_n^0}{\sqrt{z}} + O(\sqrt{z}) \quad \text{as } z \to 0,\ z \in \mathbb{C}_+, \quad c_n^0 \neq 0.
\end{aligned} \tag{2.22}$$

These facts are well known in inverse spectral theory ([N-Z], [MO], [K5]). A detailed analysis of H_0 was done in [Zh3], [K4].

If $\mu_n \in g_n \neq \emptyset$, then the function m_+ has a pole at $z_n^0 = \mu_n + i0 \in g_n^+$ (corresponding to a bound state) or at $z_n^0 = \mu_n - i0 \in g_n^-$ (corresponding to an antibound state). If $\mu_n = e_n^+$ (or $\mu_n = e_n^-$), then $z_n^0 = \mu_n$ is a virtual state. Note that for closed gaps $g_n = \emptyset, n \neq 0$, each $\psi^\pm(x,\cdot) \in \mathscr{A}(\mu_n), x \geqslant 0$. Moreover, the resolvent $R_0(z)$ has a pole at z_0 if and only if the function $m_+(\cdot)$ has a pole at z_0.

The following asymptotics from [MO] hold true as $n \to \infty$:

$$\mu_n = \pi n + \varepsilon_n(p_0 - p_{cn} + O(\varepsilon_n)), \qquad p_{cn} = \int_0^1 p(x)\cos 2\pi n x\, dx, \qquad \varepsilon_n = \frac{1}{2\pi n}, \tag{2.23}$$

$$h_{sn} = -\varepsilon_n(p_{sn} + O(\varepsilon_n)), \tag{2.24}$$

$$e_n^\pm = \pi n + \varepsilon_n(p_0 \pm |p_n| + O(\varepsilon_n)), \qquad p_n = \int_0^1 p(x)e^{-i2\pi n x}dx = p_{cn} - ip_{sn}. \tag{2.25}$$

The fundamental system y_1, y_2 of the equation $-y'' + (p+q)y = z^2 y, z \in \mathbb{C}$, with initial data

$$y_2'(t,z) = y_1(t,z) = 1, \qquad y_2(t,z) = y_1'(t,z) = 0, \tag{2.26}$$

satisfies the integral equations

$$\begin{aligned} y_1(x,z) &= \cos z(x-t) - \int_x^t \frac{\sin z(x-\tau)}{z}(p(\tau)+q(\tau))y_1(\tau,z)d\tau, \\ y_2(x,z) &= \frac{\sin z(x-t)}{z} - \int_x^t \frac{\sin z(x-\tau)}{z}(p(\tau)+q(\tau))y_2(\tau,z)d\tau. \end{aligned} \tag{2.27}$$

For each $x \in \mathbb{R}$ the functions $y_1(x,z), y_2(x,z)$ are entire in $z \in \mathbb{C}$ and satisfy

$$\begin{aligned} \max\{||z|_1 y_2(x,z)|,\ |y_2'(x,z)|, |y_1(x,z)|, \frac{1}{|z|_1}|y_1'(x,z)|\} \leqslant X_1 = e^{|\operatorname{Im} z||t-x| + \int_x^t |p+q| ds}, \\ |y_1(x,z) - \cos z(x-t)| \leqslant \frac{X_1}{|z|}\|q\|_t, \quad \left| y_2(x,z) - \frac{\sin z(x-t)}{z} \right| \leqslant \frac{X_1}{|z|^2}\|q\|_t, \end{aligned} \tag{2.28}$$

in analogy to (2.5).

The initial-value problem for the inhomogeneous equation $-y'' + (p - z^2)y = f,\ y(0) = y'(0) = 0$ has the unique solution $y(x) = -\int_0^x \varphi(x-\tau, z, \tau) f(\tau) d\tau$. Hence the solutions Φ and $\Psi^\pm$ of $-y'' + (p+q)y = z^2 y$ satisfy

$$\Phi(x,z) = \varphi(x,z) + \int_0^x \varphi(x-s, z, s) q(s) \Phi(s,z)\, ds, \tag{2.29}$$

$$\Psi^\pm(x,z) = \psi^\pm(x,z) - \int_x^t \varphi(x-s, z, s) q(s) \Psi^\pm(s,z)\, ds. \tag{2.30}$$

Below we need the well known fact from scattering theory that

$$\Psi^+(0,z) = D(z^2) = \det(I + q(H_0 - z^2)^{-1}) \ (z \in \mathcal{M}). \tag{2.31}$$

This is similar to the case $p = 0$, see [J]. The case on the real line with $p \neq$ const was considered in [F4]. The functions $k, \Psi^\pm, m_\pm, \psi^\pm$ are meromorphic in $\mathcal{Z}$ and real on $i\mathbb{R}$ and then they satisfy

$$k(-z) = \overline{k}(\overline{z}), \quad \Psi^\pm(\cdot,-z) = \overline{\Psi}^\pm(\cdot,\overline{z}), \quad m_\pm(-z) = \overline{m}_\pm(\overline{z}), \quad \psi^\pm(\cdot,-z) = \overline{\psi}^\pm(\cdot,\overline{z}) \tag{2.32}$$

for all $z \in \mathcal{Z}$.

Lemma 2.2. *i) The following identities, estimate and asymptotics hold true.*

$$\Psi^\pm = \psi^\pm(t,\cdot) y_1 + \partial_t \psi^\pm(t,\cdot) y_2, \tag{2.33}$$

$$\Psi^\pm(0,z) = 1 + \int_0^t \varphi(x,z) q(x) \Psi^\pm(x,z)\, dx, \tag{2.34}$$

$$|\Psi^\pm(x,z) - \psi^\pm(x,z)| \leqslant e^{\mp vx + b(t,x)} \frac{w^\pm(z)}{|z|_1} \int_x^t |q(r)|\, dr, \qquad (x \in [0,t]), \tag{2.35}$$

where

$$v = \operatorname{Im} k(z), \quad w^\pm = \sup_{x \in [0,t]} |e^{\pm vx} \psi^\pm(x,z)|, \quad b(t,x) = (|v| - v)(t-x) + \int_x^t (|p(r)| + |q(r)|)\, dr$$

and

$$\Psi^\pm(x,z) = e^{\pm ik(z)x}(1 + e^{\pm(t-x)(|v|-v)} O(1/z)) \tag{2.36}$$

as $|z| \to \infty, z \in \mathcal{Z}_\varepsilon$, $\varepsilon > 0$, *uniformly in* $x \in [0,t]$. *Moreover,* (1.7) *holds true.*
ii) The function $\Psi^\pm(0,\cdot)$ *has exponential type* $2t$ *in the half plane* $\mathbb{C}_\mp$.

An entire function $f(z)$ is said to be *of exponential type* if there is a constant A such that $|f(z)| \leqslant \text{const}\ e^{A|z|}$ everywhere [Koo]. The infimum over the set of A for which such an inequality holds is called the *type* of f.

Proof. i) Using (2.26), (2.14) we obtain (2.33).

The identity $\varphi(x,z,\tau) = \vartheta(\tau,z)\varphi(x+\tau,z) - \varphi(\tau,z)\vartheta(x+\tau,z)$ gives $\varphi(-x,z,x) = -\varphi(x,z)$ and (2.30) yield (2.34).

The estimate (2.35) was proved in [K4]. Substituting (2.12) into (2.35) we obtain (2.36).

In particular, substitution of (2.36), (2.28) into (2.34) yields (1.7), since $D(z^2) = \Psi^+(0,z)$.

ii) We give the proof for the case $\Psi^+(0,z)$, the proof for $\Psi^-(0,z)$ being similar. Due to (2.36), $\Psi^+(0,z)$ has exponential type $\leqslant 2t$ in the half plane $\mathbb{C}_-$. The decompositions

$$f(x,z) \equiv e^{-ixk(z)}\Psi^+(x,z) = 1 + \varrho f_1(x,z), \qquad \varphi(x,z)e^{ixk(z)} \equiv \varrho(e^{i2xz} - 1 + \varrho\eta(x,z)),$$

where $\varrho = \frac{1}{2iz}$, give

$$\begin{aligned}\Psi^+(0,z) - 1 &= \int_0^t \varphi(x,z)e^{ixk}q(x)f(x,z)\,dx \\ &= \varrho\int_0^t e^{i2xz}q(x)f(x,z)(1+\varrho e^{-i2xz}\eta(x,z))\,dx - \varrho\int_0^t q(x)f(x,z)\,dx \\ &= \varrho K(z) - \varrho\int_0^t q(x)\,dx,\end{aligned}$$

$$K = \int_0^t e^{i2xz}q(x)(1+G(x,z))\,dx, \qquad G = \varrho f_1 + \varrho e^{-i2xz}(\eta f - f_1), \qquad z \in \mathcal{Z}_\varepsilon. \tag{2.37}$$

Asymptotics (2.5), (2.36) and $k(z) = z + O(1/z)$ as $z \to \infty$ (see [KK]) yield

$$\eta(x,z) = e^{2x|\operatorname{Im} z|}O(1), \qquad f_1(x,z) = e^{2(t-x)|\operatorname{Im} z|}O(1) \quad \textit{as } |z| \to \infty,\ z \in \mathcal{Z}_\varepsilon. \tag{2.38}$$

Now we use the following variant of the Paley-Wiener Theorem from [Fr].
Let $h \in \mathcal{Q}_t$ *and let each* $F(x,z), x \in [0,t]$, *be analytic for* $z \in \mathbb{C}_-$ *and* $F \in L^2((0,t)\times\mathbb{R})$. *Then* $\int_0^t e^{2izx}h(x)(1+F(x,z))\,dx$ *has exponential type at least* $2t$ *in* $\mathbb{C}_-$.

We cannot apply this result to the function $K(z), z \in \mathbb{C}_-$, since $\psi^+(x,z)$ has a singularity at z_n^0 if $g_n \neq \emptyset$. However, we can use it for the function $K(z-i), z \in \mathbb{C}_-$, since (2.37), (2.38) imply that $\sup_{x\in[0,1]}|G(x,-i+\tau)| = O(1/\tau)$ as $\tau \to \pm\infty$. Thus the function $\Psi^+(0,z)$ has exponential type $2t$ in the half plane $\mathbb{C}_-$. ■

3. Spectral properties of H

For the sake of brevity, we write

$\varphi_1 = \varphi(1,\cdot)$, $\varphi_1' = \varphi'(1,\cdot)$, $\vartheta_1 = \vartheta(1,\cdot)$, $\vartheta_1' = \vartheta'(1,\cdot)$, $\Phi_1 = \Phi(1,\cdot)$, $\Phi_1' = \Phi'(1,\cdot)$, $\psi_t^\pm = \psi^\pm(t,\cdot)$.

We define the function

$$F(z) = \varphi(1,z)\Psi^-(0,z)\Psi^+(0,z)\ (z \in \mathcal{Z}), \tag{3.1}$$

which is real on $\mathbb{R}$, since $\Psi^-(0,z) = \overline{\Psi}^+(0,\overline{z})$ for all $z \in \mathcal{Z}$, see also (3.2).

A function f is said to belong to the *Cartwright class* $Cart_\omega$ if f is entire, of exponential type and satisfies:

$$\omega_\pm(f) := \limsup_{y\to\infty} \frac{\log|f(\pm iy)|}{y} = \omega > 0, \qquad \int_{\mathbb{R}} \frac{\log(1+|f(x)|)dx}{1+x^2} < \infty.$$

In the following, the partial derivative w.r.t. t is denoted by a dot.

Lemma 3.1. *i) The following identities and estimates hold true.*

$$F = \varphi(1,\cdot,t)y_1^2(0,\cdot) + \dot\varphi(1,\cdot,t)y_1(0,\cdot)y_2(0,\cdot) - \vartheta'(1,\cdot,t)y_2^2(0,\cdot) \in Cart_{1+2t}, \tag{3.2}$$

$$\varphi_1\dot\psi_t^+\dot\psi_t^- = -\vartheta'(1,\cdot,t), \qquad \varphi_1(\dot\psi_t^+\psi_t^- + \psi_t^+\dot\psi_t^-) = \dot\varphi(1,\cdot,t) = \varphi'(1,\cdot,t) - \vartheta(1,\cdot,t), \tag{3.3}$$

$$|F(z) - \frac{\sin z}{z}| \leqslant \frac{C_F e^{(1+2t)|\operatorname{Im} z|}}{|z|^2}, \qquad C_F = 3(\|p\|_1 + \|p+q\|_t)e^{2\|p+q\|_t + \|p\|_1}. \tag{3.4}$$

ii) The set of zeros of F *is symmetric w. r. t. both the real line and the imaginary line. In each disc* $\{z \mid |z-\pi n| < \frac{\pi}{4}\}, |n| \geqslant 1 + 4C_F e^{t\frac{\pi}{2}}$, *there exists exactly one simple real zero* z_n *of* F, *and* F *has no zeros in the domain* $\mathscr{D}_F \cap \mathbb{C}_-$, *where* $\mathscr{D}_F = \{z \in \mathbb{C} \mid 4C_F e^{2|\operatorname{Im} z|} < |z|\}$.
iii) For all $z \in \mathcal{Z}$,

$$\Psi_0^\pm(z) = e^{\pm ik(z)n_t} w_\pm(z), \quad w_\pm(z) = \Phi'(n_t, z) - m_\pm(z)\Phi(n_t, z). \tag{3.5}$$

Proof. i) The function $\varphi(1,z,t)$, $(t,z) \in \mathbb{R}\times\mathbb{C}$, (see (2.15)) satisfies [Tr]

$$\varphi(1,\cdot,t) = -\vartheta_1'\varphi_t^2 + \varphi_1\vartheta_t^2 + 2\beta\varphi_t\vartheta_t = \varphi_1\psi_t^+\psi_t^-. \tag{3.6}$$

Hence we obtain

$$\begin{aligned} &\dot\varphi(1,\cdot,t) = \varphi_1(\dot\psi_t^+\psi_t^- + \psi_t^+\dot\psi_t^-), \\ &\ddot\varphi(1,\cdot,t) = \varphi_1(\ddot\psi_t^+\psi_t^- + \psi_t^+\ddot\psi_t^- + 2\dot\psi_t^+\dot\psi_t^-) = 2\varphi_1(p(t)-z^2)\psi_t^+\psi_t^- + 2\varphi_1\dot\psi_t^+\dot\psi_t^-. \end{aligned} \tag{3.7}$$

Identity (2.33) gives

$$F = \varphi_1(\psi_t^+\psi_t^- y_1^2(0,\cdot) + \dot\psi_t^+\dot\psi_t^- y_2^2(0,\cdot) + (\psi_t^+\dot\psi_t^- + \dot\psi_t^+\psi_t^-)y_1(0,\cdot)y_2(0,\cdot). \tag{3.8}$$

Then using the following identities from [IM],

$$\dot\varphi(1,z,t) = \varphi'(1,z,t) - \vartheta(1,z,t), \quad \ddot\varphi(1,z,t) = 2(p(t)-z^2)\varphi(1,z,t) - 2\vartheta'(1,z,t), \tag{3.9}$$

and (3.6), (3.8), we obtain (3.2), (3.3), since Lemma 2.2, ii) and (2.5) yield $F \in Cart_{1+2t}$.

Next we show (3.4). We have

$$y_1(0,\cdot) = \cos tz + \widetilde{y}_1, \qquad \widetilde{y}_1 = \int_0^t \frac{\sin zs}{z}(p(s)+q(s))y_1(s,\cdot)\,ds, \tag{3.10}$$

$$y_2(0,\cdot) = -\frac{\sin tz}{z} + \widetilde{y}_2, \qquad \widetilde{y}_2 = \int_0^t \frac{\sin zs}{z}(p(s)+q(s))y_2(s,\cdot)\,ds, \tag{3.11}$$

$$\vartheta(1,t) = \cos z + \vartheta_{1t}, \qquad \vartheta_{1t} = \int_0^1 \frac{\sin z(1-s)}{z} p(s+t)\vartheta(s,t)\,ds, \tag{3.12}$$

$$\vartheta'(1,t) = -z\sin z + \vartheta_{1t}', \qquad \vartheta_{1t}' = \int_0^1 \cos z(1-s)p(s+t)\vartheta(s,t)\,ds, \tag{3.13}$$

$$\varphi(1,t) = \frac{\sin z}{z} + \varphi_{1t}, \qquad \varphi_{1t} = \int_0^1 \frac{\sin z(1-s)}{z} p(s+t)\varphi(s,t)\,ds. \tag{3.14}$$

Then (3.2) implies

$$F=(\cos tz+\widetilde{y}_1)^2(\frac{\sin z}{z}+\varphi_{1t})+(\frac{\sin tz}{z}-\widetilde{y}_2)^2(z\sin z-\vartheta_{1t}')+(\cos tz+\widetilde{y}_1)(-\frac{\sin tz}{z}+\widetilde{y}_2)\dot{\varphi}(1,z,t)$$
$$=\frac{\sin tz}{z}+f_1+f_2+f_3, \tag{3.15}$$

where

$$f_1=y_1(0,\cdot)^2\varphi_{1t}+\frac{\sin tz}{z}(\cos tz+y_1(0,\cdot))\widetilde{y}_1,$$
$$f_2=-y_2(0,\cdot)^2\vartheta_{1t}'+z\sin z(y_2(0,\cdot)-\frac{\sin tz}{z})\widetilde{y}_2, \qquad f_3=y_1(0,\cdot)y_2(0,\cdot)\dot{\varphi}(1,z,t).$$

The estimates

$$|f_3|\leqslant\frac{C_t}{|z|^2}\|p\|_1, \quad |f_j|\leqslant\frac{C_t}{|z|^2}(\|p\|_1+2\|p+q\|_t)\ (j\in\{1,2\})$$

yield (3.4), where $C_t=e^{(2t+1)|\operatorname{Im}z|+2\|p+q\|_t+\|p\|_1}$.

ii) From (3.4) we obtain for $|n|\geqslant 1+4C_Fe^{t\frac{\pi}{2}}$

$$|F(z)-\frac{\sin z}{z}|\leqslant\frac{C_F}{|z|^2}e^{|\operatorname{Im}z|+t\frac{\pi}{2}|}\leqslant\frac{4C_F}{|z|}e^{t\frac{\pi}{2}|}\frac{|\sin z|}{|z|}<\frac{|\sin z|}{|z|}$$

for all $|z-\pi n|=\frac{\pi}{4}$, since $e^{|\operatorname{Im}z|}\leqslant 4|\sin z|$ for all $|z-\pi n|\geqslant\pi/4, n\in\mathbb{Z}$ (see [PT]). Hence, by Rouché's theorem, F has the same number of roots, counting multiplicities, as $\sin z$ in each disc $\mathcal{D}_{\frac{\pi}{4}}(\pi n)$. Since $\sin z$ only has the roots πn ($n\in\mathbb{Z}$), the statement about the zeros of F in each disc follows. The zero in $\mathcal{D}_{\frac{\pi}{4}}(\pi n)$ is real, since F is real-valued on the real line.

Using (3.4) and $e^{|\operatorname{Im}z|}\leqslant 4|\sin z|$ for all $|z-\pi n|\geqslant\pi/4, n\in\mathbb{Z}$, we obtain

$$|F(z)|\geqslant|\frac{\sin z}{z}|-\left|F(z)-\frac{\sin z}{z}\right|\geqslant\frac{e^{|\operatorname{Im}z|}}{4|z|^2}\left(|z|-4C_Fe^{2t|\operatorname{Im}z|}\right)>0$$

for all $z\in\mathscr{D}_1=\{z\in\mathscr{D}_Fz \mid |z-\pi n|\geqslant\pi/4(n\in\mathbb{Z})\}$. As the function F has exactly one real zero z_n in $\mathcal{D}_{\frac{\pi}{4}}(\pi n), n\geqslant n_0$, it therefore has no zeros in the domain $\mathscr{D}_F$. That the set of zeros of F is symmetric with respect to both the real line and the imaginary line follows from the fact that F is real-valued on both lines.

iii) Using $\Psi^{\pm}(0,z)=\{\Psi^{\pm}(x,z),\Phi(x,z)\}$ ($z\in\mathcal{Z}$) at $x=n_t$, and (2.11) we obtain (3.5), where $\{y,u\}=yu'-y'u$ is the Wronskian. ■

With $g_n^c=\overline{g}_n^-\cup\overline{g}_n^+$ as before, we define the sets

$$\mathscr{X}=i\mathbb{R}\cup\mathbb{C}_-\cup\bigcup_{n\in\mathbb{Z}}g_n^c\subset\mathcal{Z}, \qquad \mathscr{X}_0=\mathscr{X}\setminus\{0,e_n^{\pm},\mu_n\pm i0(n\in\mathbb{Z},g_n\neq\emptyset)\}. \tag{3.16}$$

We describe the zeros of $\Psi^{\pm}(0,z)$.

Lemma 3.2. *i) If $g_n=(e_n^-,e_n^+)=\emptyset$ for some $n\neq 0$, then $\Psi^{\pm}(x,\cdot)\in\mathscr{A}(\mu_n)$ $(x\geqslant 0)$, and* $\operatorname{Im}\Psi^{\pm}(0,\mu_n)\neq 0$. *Moreover, $\mu_n=e_n^{\pm}$ is a simple zero of F and $\mu_n\notin\sigma_{st}(H)$.*
ii) $\Psi^{\pm}(0,z)\neq 0$ for all $z\in(e_{n-1}^+,e_n^-), n\in\mathbb{Z}$. Moreover, states of H and zeros of $\Psi^+(0,z)$ lie in $\mathscr{X}$.
iii) A point $z\in\mathscr{X}_0$ is a zero of Ψ_0^+ if and only if $z\in\mathscr{X}_0\cap\mathfrak{S}_{st}(H)$. In particular, if $z\in\mathscr{X}_0\cap\mathfrak{S}_{st}(H)$, then

1) $z\in\overline{\mathbb{C}}_+\cap\mathcal{Z}$ is a bound state of H, *2) $z \in \overline{\mathbb{C}}_- \cap \mathcal{Z}$ is a resonance of H.*

Proof. i) Lemma 2.1 and identity (2.17) yield that $\Psi^\pm(x,\cdot) \in \mathscr{A}(\mu_n)$ $(x \geqslant 0)$.
Using (2.9) we deduce that $\Psi_0^\pm(\mu_n) \neq 0$, hence $\Psi^\pm(x,\cdot), 1/\Psi_0^\pm \in \mathscr{A}(\mu_n)$. Thus μ_n is not a state of H and μ_n is a simple zero of F.
ii) The conformal mapping k maps each interval $(e_{n-1}^+, e_n^-), n \geqslant 1$, onto $(\pi(n-1), \pi n)$. As $\mu_n \in [e_n^-, e_n^+]$, the function $m_\pm$ is analytic on (e_{n-1}^+, e_n^-) and $\operatorname{Im} m_\pm(z) \neq 0$ for all $z \in (e_{n-1}^+, e_n^-)$. Then the identity (2.17) gives $\Psi^\pm(0,z) \neq 0$ for all $z \in (e_{n-1}^+, e_n^-)$.
The identity $\Psi^+(0,z) = D(z^2)$ and standard arguments (similar to the case $p=0$, see [K1]) imply that states of H and zeros of $\Psi^+(0,z)$ belong to the set $\mathscr{Z}$.
iii) follows from the identities (2.17), (2.18). ■

States of H which coincide with unperturbed states z_n^0 have the following properties.

Lemma 3.3. *Let $\zeta = \mu_n + i0 \in g_n^+$ or $\zeta = \mu_n - i0 \in g_n^-$ for some $n \geqslant 1$, where $g_n \neq \emptyset$. Then*
i) $\widetilde{\varphi}(0,\mu_n) = 0$ if and only if $\Phi(n_t,\mu_n) = 0$.
ii) Assume in addition $\zeta = z_n^0 \in \sigma_{st}(H_0)$. Then $\Psi_0^- \in \mathscr{A}(\zeta)$, each $\Psi^+(x,\cdot)$, $x > 0$, has a simple pole at ζ, and there are two cases:
1) if $\widetilde{\varphi}(0,\mu_n) = 0$, then $\Psi_0^+ \in \mathscr{A}(\zeta)$, $\Psi_0^-(\zeta) \neq 0$ and $\zeta \in \sigma_{st}(H)$. In particular,

$$\zeta = \mu_n + i0 \in g_n^+ \quad \Rightarrow \quad \Psi_0^+(\zeta) \neq 0, \quad F(\mu_n) = 0, \qquad (-1)^n F'(\mu_n) > 0; \tag{3.17}$$

2) if $\widetilde{\varphi}(0,\mu_n) \neq 0$, then

$$\frac{\Psi^+(x,\cdot)}{\Psi_0^+} \in \mathscr{A}(\zeta) \ (x \geqslant 0), \qquad \zeta \notin \mathfrak{S}_{st}(H), \quad \Psi_0^-(\zeta) \neq 0, \quad F(\mu_n) \neq 0,$$

$$\Psi_0^+(z) = \frac{c_n + O(\varepsilon)}{\varepsilon} \widetilde{\varphi}(0) \quad \text{as } \varepsilon = z - \zeta \to 0. \tag{3.18}$$

iii) $\zeta \in \mathfrak{S}_{bs}(H)$ (or $\zeta \in \sigma_{vs}(H)$) if and only if $\zeta \in \mathfrak{S}_{bs}(H_0)$ (or $\zeta \in \sigma_{vs}(H_0)$) and $\Phi(n_t,\mu_n) = 0$.
iv) If $\zeta \in \mathfrak{S}_{st}(H_0) \cap \mathfrak{S}_{st}(H)$, then its complex conjugate $\overline{\zeta}$ on the other sheet is not a state of H.
v) Let $\zeta \in \mathfrak{S}_{st}(H_0)$ and assume that its complex conjugate $\overline{\zeta}$ on the other sheet is a state of H. Then $\zeta \notin \mathfrak{S}_{st}(H)$.

Proof. i) Comparing (3.5) and (2.17) we deduce that $\widetilde{\varphi}(0,\mu_n) = 0$ if and only if $\Phi(0,\mu_n) = 0$.
ii) Lemma 2.1 yields $m_- \in \mathscr{A}(\zeta)$ and each $\Psi^+(x,\cdot)$, $x > 0$, has a simple pole at ζ and $m_+(z) = \frac{\pm c_n}{\varepsilon} + O(1)$ as $\varepsilon \to 0$, $\varepsilon \in \mathbb{C}_\pm$, with $c_n < 0$.
1) If $\widetilde{\varphi}(0,\mu_n) = 0$, then (2.17) yields $\Psi_0^+ \in \mathscr{A}(\zeta)$, $\Psi_0^-(\zeta) \neq 0$. Thus $\zeta \in \mathfrak{S}_{st}(H)$, since each $\Psi^+(x,\cdot)$, $x > 0$¡ has a simple pole at ζ.
Consider the case $\zeta = \mu_n + i0 \in g_n^+$. As (3.5) gives

$$\Psi^\pm(0,z) = e^{\pm ik(z)n_t} w_\pm(z), \quad w_\pm(z) = \Phi'(n_t,z) - m_\pm(z)\Phi(n_t,z), \qquad (z \in \mathcal{Z}),$$

we find

$$w_-(\zeta) = \Phi'(n_t,\mu_n) \neq 0, \quad w_+(z_1) = \Phi'(n_t,\mu_n)(1 - c_n \frac{\partial_z \Phi(n_t,\mu_n)}{\Phi'(n_t,\mu_n)}) \neq 0,$$

since $\Phi'(n_t,\mu_n) \neq 0, \Phi(n_t,\mu_n) = 0$ and $\Phi'(n_t,\mu_n)\partial_z\Phi(n_t,\mu_n) > 0$ [PT]. This yields (3.17), since $(-1)^n \partial_z \varphi(1,\mu_n) > 0$ [PT].

2) Lemma 2.1 gives $m_+(z) = \frac{c_n+O(\varepsilon)}{\varepsilon}$ as $\varepsilon = z-\zeta \to 0$. Using (2.17), we obtain

$$\Psi_0^+(z) = \frac{c_n + O(\varepsilon)}{\varepsilon}\widetilde{\varphi}(0), \qquad \frac{\Psi^+(x,z)}{\Psi_0^+(z)} = \frac{\varepsilon\widetilde{\vartheta}(x) - c_n\widetilde{\varphi}(x) + O(\varepsilon)}{\varepsilon\widetilde{\vartheta}(0) - c_n\widetilde{\varphi}(0) + O(\varepsilon)} = \frac{\widetilde{\varphi}(x)}{\widetilde{\varphi}(0)} + O(\varepsilon) \quad \text{as } \varepsilon \to 0,$$

since $c_n\widetilde{\varphi}(0,\mu_n) \neq 0$, where we abbreviate $\widetilde{\varphi}(x) = \widetilde{\varphi}(x,z)$ etc. This yields $\varphi(1,z)\Psi_0^+(z) = \partial_z\varphi(1,\mu_n)c_n + o(1)$, and $m_- \in \mathscr{A}(\zeta)$ gives $\Psi_0^+(\zeta) = \widetilde{\vartheta}(0) \neq 0$, so $F(\mu_n) \neq 0$ and (3.18).

Using i) and ii) we obtain iii).

iv) $\Psi_0^+ \in \mathscr{A}(\overline{\zeta})$ and each $\Psi^+(x,\cdot) \in \mathscr{A}(\overline{\zeta}), x > 0$. Due to ii) $\widetilde{\varphi}(0,\mu_m) = 0$, then we obtain $\Psi_0^+(\overline{\zeta}) \neq 0$. Thus $\overline{\zeta} \notin \mathfrak{S}_{st}(H)$.

v) follows directly from iv) by contraposition. ■

For virtual states, which coincide with the points $e_n^\pm$, we have the following.

Lemma 3.4. *Let* $\zeta = e_n^-$ *or* $\zeta = e_n^+$ *for some* $n \geqslant 1$, *where* $e_n^- < e_n^+$ *and set* $\varepsilon = z - \zeta$.
i) Assume that $\zeta \neq \mu_n$ *and* $\Psi_0^+(\zeta) = 0$. *Then* ζ *is a simple zero of* F, $\zeta \in \sigma_{vs}(H)$ *and*

$$\Psi_0^+(z) = \widetilde{\varphi}(0,\zeta)c\sqrt{\varepsilon} + O(\varepsilon), \quad \mathscr{R}(x,z) = \frac{\Psi^+(x,z) + O(\varepsilon)}{\widetilde{\varphi}(0,\zeta)c\sqrt{\varepsilon}}, \quad c\widetilde{\varphi}(0,\zeta) \neq 0. \tag{3.19}$$

ii) Assume that $\zeta = \mu_n$ *and* $\widetilde{\varphi}(0,\zeta) \neq 0$. *Then* $F(\zeta) \neq 0$ *and each* $\mathscr{R}(x,\cdot), x > 0$, *does not have a singularity at* ζ, *and* $\zeta \notin \sigma_{vs}(H)$.
iii) Assume that $\zeta = \mu_n$ *and* $\widetilde{\varphi}(0,\zeta) = 0$. *Then* $\zeta \in \sigma_{vs}(H)$, $\Psi_0^\pm(\zeta) \neq 0$ *and* ζ *is a simple zero of* F, *and each* $\mathscr{R}^2(x,\cdot), x > 0$, *has a pole at* ζ.

Proof. i) Lemma 2.1 gives $m_\pm(z) = m_\pm(\zeta) + c\sqrt{\varepsilon} + O(\varepsilon)$ as $\varepsilon = z - \zeta \to 0, c \neq 0$. We distinguish two cases. First assume $\widetilde{\varphi}(0,\zeta) \neq 0$. Then the identity (2.17) implies (3.19).

Second, if $\widetilde{\varphi}(0,\zeta) = 0$, then (2.17) implies $\Psi_0^+(\zeta) = \widetilde{\vartheta}(0,\zeta) \neq 0$, which gives a contradiction.
ii) If $\zeta = \mu_n$, then Lemma 2.1 gives $m_\pm(z) = \pm\frac{c}{\sqrt{\varepsilon}} + O(1)$ $(\varepsilon \to 0)$ with $c \neq 0$. Then (2.17) implies

$$\Psi_0^\pm(z) = \pm\frac{\widetilde{\varphi}(0,\zeta)c}{\sqrt{\varepsilon}} + O(1), \qquad \frac{\Psi^+(x,z)}{\Psi_0^+(z)} = \frac{\widetilde{\vartheta}(x,z) + (\frac{c}{\sqrt{\varepsilon}} + O(1))\widetilde{\varphi}(x,z)}{\frac{\widetilde{\varphi}(0,\zeta)c}{\sqrt{\varepsilon}} + O(1)} = \frac{1 + O(\sqrt{\varepsilon})}{\widetilde{\varphi}(0,\zeta)}.$$

Thus the function $\mathscr{R}(x,\cdot), x > 0$, is not singular at ζ, and $\zeta \notin \sigma_{vs}(H)$, $F(\zeta) \neq 0$.

iii) If $\widetilde{\varphi}(0,\zeta) = 0$, then (2.17) gives $\Psi_0^+(\zeta) = \widetilde{\vartheta}(0,\zeta) \neq 0$, since $\widetilde{\vartheta}(0,\zeta) \neq 0$ and $\beta(\zeta) = 0$. Moreover, we obtain $\Psi^+(x,z) = \widetilde{\vartheta}(x,z) + (\frac{c}{\sqrt{\varepsilon}} + O(1))\widetilde{\varphi}(x,z)$, and the function $\mathscr{R}^2(x,\cdot), x > 0$, has a pole at ζ, $\zeta \in \sigma_{vs}(H)$ and $F(\zeta) = 0$. ■

Lemma 3.5. *Let* $\lambda \in \gamma_n$, $\lambda \neq \mu_n^2$, *be an eigenvalue of* H *for some* $n \geqslant 0$ *and let* $z = \sqrt{\lambda} \in i\mathbb{R}_+ \cup \bigcup_{n\geqslant 1} g_n^+$. *Then*

$$C_\lambda = \int_0^\infty |\Psi^+(x,z)|^2\,dx = -\frac{\Psi^{+\prime}(0,z)}{2z}\partial_z\Psi^+(0,z) > 0, \tag{3.20}$$

$$\frac{2i\sin k(z)}{\varphi(1,z)} = \Psi^-(0,z_1)\Psi^{+\prime}(0,z) \neq 0, \qquad i\sin k(z) = -(-1)^n\sinh h, \quad h = h(\lambda) > 0, \tag{3.21}$$

$$C_\lambda = \frac{(-1)^n F'(z)\sinh h}{z\varphi^2(1,z)\Psi^-(0,z)^2} > 0, \qquad \frac{(-1)^n F'(z)}{z} > 0. \tag{3.22}$$

Proof. (3.20) follows from the identity $\{\frac{\partial}{\partial z}\Psi^+, \Psi^+\}' = 2z(\Psi^+)^2$.

Using the Wronskian of the functions Ψ^+, Ψ^- and (2.11), we obtain $\Psi^-(0,z)\Psi^{+\prime}(0,z) = m_+(z) - m_-(z)$, which yields (3.21), since $k(z) = \pi n + ih$ for some $h > 0$ (see the definition of k before (2.11)). Then the identities (3.20), (3.21) imply (3.22). ■

4. Proof of the Main Theorems

Proof of Theorem 1.1. i) The asymptotics (1.7) were proved in Lemma 2.2.

ii) and iii) of Lemma 3.3 give (1.9) for the case of non-virtual states, i.e., those not coinciding with $e_n^\pm$.

Lemma 3.4 implies (1.9) for the case of virtual states.

Lemma 3.2 gives (1.8) for the case of non-virtual states. Lemma 3.4 implies (1.8) for the case of virtual states.

ii) Using ii) and iii) of Lemma 3.3 we obtain (1.10).

iii) Due to i) ζ is a zero of F, so (3.4) yields (1.11). Lemma 3.1 ii) completes the proof of iii). ■

Proof of Theorem 1.2. i) Let $g_n \neq \emptyset$. The entire function $F = \varphi(1,\cdot)\Psi_0^+\Psi_0^-$ has opposite sign on σ_n and σ_{n+1}, since $\Psi_0^+(z)\Psi_0^-(z) = |\Psi_0^+(z)|^2 > 0$ for z inside $\sigma_n \cup \sigma_{n+1}$ (see (2.33)) and $\varphi(1,\cdot)$ has one simple zero in each interval $[e_n^-, e_n^+]$. Therefore F has an odd number of zeros on $[e_n^-, e_n^+]$.

By Lemma 3.2-3.4, $\zeta \in g_n^c$ is a state of H if and only if $\zeta \in \overline{g}_n$ is a zero of F (with equal multiplicities). Hence the number of states (counting multiplicities) on g_n^c is odd.

Using Lemma 3.1 and 3.2, we deduce that there exists exactly one simple state z_n in each interval $[e_n^-, e_n^+]$ for $g_n \neq \emptyset$ and for sufficiently large $n \geqslant 1$. Moreover, the asymptotics (2.25) $e_n^\pm = \pi n + \frac{p_0}{2\pi n} + o(1/n)$ give

$$z_n = \pi n + \frac{p_0}{2\pi n} + o(1/n). \tag{4.1}$$

By arguments analogous to the proof of (2.34), we obtain the identities

$$\widetilde{\vartheta}(0,z) = 1 + \int_0^t \varphi(x,z)q(x)\widetilde{\vartheta}(x,z)dx, \qquad \widetilde{\varphi}(0,z) = \int_0^t \varphi(x,z)q(x)\widetilde{\varphi}(x,z)dx; \tag{4.2}$$

then a standard iteration procedure and (4.1) give the asymptotics

$$\widetilde{\vartheta}(0,z_n) = 1 + O(1/n), \tag{4.3}$$

$$\widetilde{\varphi}(0,z_n) = \int_0^t \frac{\sin^2 z_n x}{z_n^2} q(x)\,dx + O(1/n^3) = \frac{q_0 - \widehat{q}_{cn}}{2(\pi n)^2} + O(1/n^3), \tag{4.4}$$

where $\widehat{q}_{cn} = \int_0^t q(x)\cos 2\pi n x\,dx$. Using (2.13) and $\Psi^\pm = \widetilde{\vartheta} + m_\pm\widetilde{\varphi}$, see (2.17), we obtain

$$F = F_1 + F_2 + F_3, \qquad F_1 = \varphi(1,\cdot)\widetilde{\vartheta}_0^2, \quad F_2 = 2\beta\widetilde{\vartheta}_0\widetilde{\varphi}_0, \qquad F_3 = -\vartheta'(1,\cdot)\widetilde{\varphi}_0^2, \tag{4.5}$$

where we abbreviate $\widetilde{\vartheta}_0 = \widetilde{\vartheta}(0,z)$, $\widetilde{\varphi}_0 = \widetilde{\varphi}(0,z)$. Using the estimates (4.3), (4.4), we obtain

$$\begin{gathered} F_1(z_n) = \varphi(1,z_n)(1+O(1/n)), \qquad\qquad F_3(z_n) = O(1/n^4) \\ F_2(z_n) = (-1)^n \frac{(p_{sn} + O(1/n))}{\pi n}\frac{(q_0 - \widehat{q}_{cn} + O(1/n))}{2(\pi n)^2} = f_n + O(1/n^4) \quad \text{as } n \to \infty, \end{gathered} \tag{4.6}$$

where $f_n = (-1)^n \frac{p_{sn}(q_0-\widehat{q}_{cn})}{2(\pi n)^3}$. Combining these asymptotics with $F(z_n)=0$, we get

$$\varphi(1,z_n) = -F_2(z_n) + O(1/n^4) = -f_n + O(1/n^4). \tag{4.7}$$

Then, using $\varphi(1,z_n) = \partial_z\varphi(1,\mu_n)\delta_n + O(1/n^4)$, where $z_n = \mu_n + \delta_n$, we obtain

$$\partial_z\varphi(1,\mu_n)\delta_n = -f_n + O(1/n^4)$$

and the asymptotics $\partial_z\varphi(1,\mu_n) = \frac{(-1)^n}{(\pi n)} + O(1/n^2)$ give

$$\delta_n = -\frac{f_n}{\partial_z\varphi(1,\mu_n)} + O(1/n^3) = -\frac{(\widehat{q}_0-\widehat{q}_{cn})p_{sn}}{2(\pi n)^2} + O(1/n^3),$$

which yields (1.14).

Now we denote by $\mathcal{N}(r,f)$ the total number of zeros of f of modulus $\leqslant r$, and by $\mathcal{N}^+(r,f)$ the number of zeros counted in $\mathcal{N}(r,f)$ with non-negative real part, by $\mathcal{N}^-(r,f)$ those with negative real part (each zero being counted according to its multiplicity). Then the following result holds [Koo].

Theorem (Levinson). *Let the entire function $f\in Cart_\omega$. Then $\mathcal{N}^\pm(r,f) = \frac{r}{\pi}(\omega+o(1))$ as $r\to\infty$, and for each $\delta>0$ the number of zeros of f of modulus $\leqslant r$ lying outside both of the two sectors $|\arg z|<\delta$, $|\arg z-\pi|<\delta$ is $o(r)$ for $r\to\infty$.*

We also denote by $\mathcal{N}_+(r,f)$ (or $\mathcal{N}_-(r,f)$) the number of zeros of f counted in $\mathcal{N}(r,f)$ with non-negative (or negative) imaginary part, each zero being counted according to its multiplicity.

Let $s_0=0$ and $\pm s_n>0, n\in\mathbb{N}$, be all real zeros of F and let n_0 be the multiplicity of the zero $s_0=0$. Consider the entire function $F_1 = z^{n_0}\lim_{r\to\infty}\prod_{0<s_n\leqslant r}(1-\frac{z^2}{s_n^2})$. The Levinson Theorem and Lemma 2.2 imply

$$\mathcal{N}(r,F) = \mathcal{N}(r,F_1) + \mathcal{N}(r,F/F_1) = 2r\frac{1+2t}{\pi} + o(r), \quad \mathcal{N}(r,F_1) = \frac{2r}{\pi} + o(r), \tag{4.8}$$

$$\mathcal{N}_-(r,F) = \mathcal{N}_+(r,F) = \mathcal{N}_-(r,\Psi_0^+) - N_0, \tag{4.9}$$

as $r\to\infty$, where N_0 is the number of non-positive eigenvalues of H. Thus

$$2\mathcal{N}_-(r,F) = 2r\frac{2t}{\pi} + o(r), \tag{4.10}$$

which yields (1.15).

ii) Using Lemma 3.5 we obtain the statements ii) and iii). ■

Proof of Theorem 1.3. Let $z=e_n^\pm$. Identity (3.5) and $k(e_n^\pm)=\pi n$ yield

$$\Psi_0^-(z) = \Psi_0^+(z) = (-1)^N w_+(z), \quad w_+(z) = \Phi'(n_t,z) - \frac{\beta(z)}{\varphi(1,z)}\Phi(n_t,z), \quad N = n_t n. \tag{4.11}$$

Estimates (2.5) and $e_n^\pm = \pi n + \varepsilon_n(p_0 \pm |p_n| + O(\varepsilon_n))$, $\varepsilon_n = \frac{1}{2\pi n}$ (see (2.25)), give

$$\Phi'(n_t,z) = (-1)^N + O(1/n), \quad \Phi(n_t,z) = \frac{\sin n_t z}{\pi n} + O(1/n^2) = O(1/n^2). \tag{4.12}$$

Using (2.28), we obtain

$$\sin e_n^\pm = (-1)^n \sin\frac{\pm|p_n|}{2\pi n} + O(1/n^2) = (-1)^n\frac{\pm|p_n|}{2\pi n} + O(1/n^2),$$

$$\varphi(1,e_n^{\pm}) = \frac{\sin e_n^{\pm}}{\pi n} + \frac{(-1)^n p_{cn}}{2\pi^2 n^2} + O(1/n^3) = (-1)^n \frac{\pm|p_n| + p_{cn}}{2\pi^2 n^2} + O(1/n^3).$$

Then the estimate $\sqrt{x^2+y^2} - y \geqslant x$ for $y, x \geqslant 0$ gives $|p_n| \pm p_{cn} \geqslant |p_{sn}|$, which yields

$$\frac{\beta(e_n^+)}{\varphi(1,e_n^+)} = \pi n \frac{p_{sn} + O(1/n)}{\pm|p_n| + p_{cn} + O(1/n)} = O(\pi n), \tag{4.13}$$

since $|p_{sn}| \geqslant \frac{1}{n^\alpha}$. Combining (4.11)-(4.13) and (2.7), we obtain $\Psi_0^+(e_n^{\pm}) = 1 + o(1)$. The function $\Psi_0^+(z)$ is analytic on g_n^- and $\Psi_0^+(e_n^{\pm}) = 1 + o(1)$. Thus $\Psi_0^+(z)$ has no zeros on g_n^-, since by Theorem 1.2, the function F has exactly one zero on each non-empty $\overline{g}_n$ for large $n > 1$.

Let $\mu_n + i0 \in g_n^+$ be a bound state of H_0 for some sufficiently large n. Then Lemma 2.1 implies $h_{sn} > 0$. Moreover, (2.24) gives $h_{sn} = -\frac{p_{sn} + O(\frac{1}{n})}{2\pi n}$ as $n \to \infty$. Thus $p_{sn} < -\frac{1}{n^\alpha}$ for large $n > 1$ and, by the asymptotics (1.14), the bound state $z_n > \mu_n$ if $q_0 > 0$ and $z_n < \mu_n$ if $q_0 < 0$. The proof of the other cases is similar. ■

Proof of Theorem 1.4. i) Using the identities (2.34) and (2.17) we obtain

$$\Psi_0^+ = Y_1 + \frac{i \sin k}{\varphi_1} \widetilde{\varphi}(z_n), \quad Y_1 = \widetilde{\vartheta}(z_n) + \frac{\beta}{\varphi_1} \widetilde{\varphi}(z_n). \tag{4.14}$$

Note that (2.13) gives $\beta(\mu_n) = 0$. Then the asymptotics (1.14), (2.6), (2.5) imply

$$\frac{\beta(z_n)}{\varphi(1,z_n)} = \frac{\beta'(\mu_n) + O(\varepsilon_n^3)}{\partial_z \varphi(1,\mu_n) + O(\varepsilon_n^3)} = o(1) \qquad \text{as} \quad n \to \infty, \qquad \varepsilon_n = \frac{1}{2\pi n} \tag{4.15}$$

where we used the asymptotics $\partial_z \varphi(1,\mu_n) = \frac{(-1)^n + O(\varepsilon_n)}{(\pi n)}$ and $\beta'(\mu_n) = o(1/n)$. Thus (4.3), (4.4) give

$$Y_1 = 1 + O(\varepsilon_n), \qquad \widetilde{\varphi}(0,z_n) = 2\varepsilon_n^2 (b_n + O(\varepsilon_n)), \qquad b_n = q_0 - \widehat{q}_{cn} \tag{4.16}$$

In the following we need these identities and asymptotics as $n \to \infty$ from [KK]:

$$(-1)^{n+1} i \sin k(z) = \sinh v(z) = \pm|\Delta^2(z) - 1|^{\frac{1}{2}} > 0 \quad (z \in g_n^{\pm}), \tag{4.17}$$

$$v(z) = \pm|(z - e_n^-)(e_n^+ - z)|^{\frac{1}{2}}(1 + O(1/n^2)), \quad \sinh v(z) = v(z)(1 + O(|g_n|^2)), \quad (z \in \overline{g}_n^{\pm}). \tag{4.18}$$

We rewrite the equation $\Psi_0^+ = 0$ in the form $\varphi_1 Y_1 = -i \sin k \widetilde{\varphi}(z_n)$. Then we obtain

$$2\delta\varepsilon_n(1 + O(\varepsilon_n)) = v(z) 2\varepsilon_n^2 (b_n + O(\varepsilon_n)) = \sqrt{\delta(|g_n| - \delta)} 2\varepsilon_n^2 (b_n + O(\varepsilon_n)),$$

$$\sqrt{\delta} = \varepsilon_n \sqrt{|g_n| - \delta} (b_n + O(\varepsilon_n)), \qquad \delta = z_n - \mu_n,$$

where $\sqrt{\delta} > 0$ if $b_n > 0$ and $\sqrt{\delta} < 0$ if $b_n < 0$. The last asymptotics imply $\delta = \varepsilon_n^2 |g_n| (b_n + O(\varepsilon_n))^2$, where $b_n = q_0 - \widehat{q}_{cn}$, which yields (1.17). ■

Proof of Theorem 1.5. i) Let $q \in \mathcal{Q}_t, q_0 = 0$ and let each $|\widehat{q}_{cn}| > n^{-\alpha}$ for some $\alpha \in (0,1)$ and for sufficiently large n. (The proof of the other cases is similar.) Using the inverse spectral theory from [K5] summarised in the Introduction above, we see that for any sequence $\varkappa = (\varkappa_n)_1^\infty \in \ell^2, \varkappa_n \geqslant 0$, there exists a potential $p \in L^2(0,1)$ such that gap γ_n has length $|\gamma_n| = \varkappa_n$ $(n \in \mathbb{N})$. Moreover, for large n we can write the gap in the form $\gamma_n = (E_n^-, E_n^+)$, where $\mu_n^2 = E_n^-$ or $\mu_n^2 = E_n^+$. We choose $\mu_n^2 = E_n^-$ or $\mu_n^2 = E_n^+$ depending on the given sequence $\sigma = (\sigma_n)_1^\infty$, where $\sigma_n \in \{0,1\}$, as follows, using Theorem 1.3 (i).

If $\sigma_n = 1$ and $\widehat{q}_{cn} < -n^{-\alpha}$ (or $\widehat{q}_{cn} > n^{-\alpha}$), then taking $\mu_n^2 = E_n^-$ (or $\mu_n^2 = E_n^+$) we ensure that λ_n is an eigenvalue for large n.

If $\sigma_n = 0$ and $\widehat{q}_{cn} > n^{-\alpha}$ (or $\widehat{q}_{cn} < -n^{-\alpha}$), then taking $\mu_n^2 = E_n^-$ (or $\mu_n^2 = E_n^+$) we ensure that λ_n is an antibound state for large n.

ii) Let $p \in L^2(0,1)$ such that $\mu_n^2 = E_n^-$ or $\mu_n^2 = E_n^+$ for sufficiently large $n \in \mathbb{N}_0$, $\gamma_n = (E_n^-, E_n^+)$. Let $\sigma = (\sigma_n)_1^\infty$ be any sequence, where $\sigma_n \in \{0,1\}$. We take $|\widehat{q}_{cn}| > n^{-\alpha}$ for large $n \in \mathbb{N}_0$, and then choose the sign of q_{cn} as follows, using Theorem 1.3 i).

If $\sigma_n = 0$ and $\lambda_n^0 = E_n^-$ (or $\lambda_n^0 = E_n^+$), then taking $\widehat{q}_{cn} > n^{-\alpha}$ (or $\widehat{q}_{cn} < -n^{-\alpha}$) we ensure that λ_n is an antibound state.

If $\sigma_n = 1$ and $\lambda_n^0 = E_n^-$ (or $\lambda_n^0 = E_n^-$), then taking $\widehat{q}_{cn} < -n^{-\alpha}$ (or $\widehat{q}_{cn} > n^{-\alpha}$) we ensure that λ_n is an eigenvalue. ■

The proof of Theorem 1.6 will be based on the following asymptotic result on the number of eigenvalues introduced in a gap by a compactly supported perturbation. This follows the original idea of [So] in the modified version of [Sc].

Lemma 4.1. *Let $q \in \mathcal{Q}_t$ be continuous and H_τ, $\Omega = [E_1, E_2]$ as in Corollary 1.6. Then*

$$\#_{bs}(H_\tau, \Omega) = \tau \int_0^\infty \Big(\varrho(E_2 - q(x)) - \varrho(E_1 - q(x)) \Big) dx + o(\tau) \quad \text{as} \quad \tau \to \infty, \tag{4.19}$$

where ϱ is the integrated density of states (1.18).

Proof. By the Glazman decomposition principle and the properties of the periodic problem on a half-line summarised in the Introduction above, the number of bound states of H_τ in Ω differs by no more than 5 from the number of eigenvalues in Ω of the regular Sturm-Liouville operator $\tilde{H}_\tau$ on $[0, t\tau]$ with Dirichlet boundary conditions. This number can be estimated by oscillation theory as follows. For $\lambda \in \Omega$, let u be the (real-valued) solution of the equation

$$-u'' + p(x)u + q(x/\tau)u = \lambda u$$

with $u(0) = 0$, $u'(0) = 1$, and $\theta(\cdot, \lambda) : [0, t\tau] \to \mathbb{R}$ the locally absolutely continuous function (called a *Prüfer angle*) with the properties

$$\begin{pmatrix} u \\ u' \end{pmatrix} = \sqrt{u^2 + u'^2} \begin{pmatrix} \sin\theta \\ \cos\theta \end{pmatrix}, \qquad \theta(0, \lambda) = 0.$$

Then $\theta(t\tau, \lambda)$ is continuous and monotone increasing as a function of λ, and the number of eigenvalues of $\tilde{H}_\tau$ in Ω differs by no more than 1 from $\frac{1}{\pi}(\theta(t\tau, E_2) - \theta(t\tau, E_1))$.

In the following we shall use the fact, which can be shown along the lines of [E2] Theorem 3.1.2, [Sc] Corollary 1, that the Prüfer angle $\tilde{\theta}$ of any real-valued solution of the periodic equation

$$-y'' + py = \lambda y$$

satisfies $\tilde{\theta}(x) = \pi\varrho(\lambda)x + O(1)$ as $x \to \infty$.

Consider a division of the interval $[0, t]$ into n parts, $0 = s_0 < s_1 < \cdots < s_{n-1} < s_n = t$. We set $q_j^+ := \sup\limits_{[s_{j-1}, s_j]} q$, $q_j^- := \inf\limits_{[s_{j-1}, s_j]} q$. Then if $\theta_j^\pm$ is the Prüfer angle of a non-trivial real-valued solution of

$$-y'' + p(x)y + q_j^\pm y = \lambda y$$

on the interval $[\tau s_{j-1}, \tau s_j]$ with $\theta_j^\pm(\tau s_{j-1}) = \theta(\tau s_{j-1})$, then Sturm comparison ([W] Thm 13.1) shows that

$$\theta_j^+(\tau s_j) \leqslant \theta(\tau s_j) \leqslant \theta_j^-(\tau s_j).$$

Hence

$$\sum_{j=1}^{n}\varrho(\lambda-q_j^+)(s_j-s_{j-1})=\lim_{\tau\to\infty}\sum_{j=1}^{n}\frac{\theta_j^+(\tau s_j,\lambda)-\theta_j^+(\tau s_{j-1},\lambda)}{\tau\pi}\leqslant\lim_{\tau\to\infty}\sum_{j=1}^{n}\frac{\theta(\tau s_j,\lambda)-\theta(\tau s_{j-1},\lambda)}{\tau\pi}$$
$$\leqslant\lim_{\tau\to\infty}\sum_{j=1}^{n}\frac{\theta_j^-(\tau s_j,\lambda)-\theta_j^-(\tau s_{j-1},\lambda)}{\tau\pi}=\sum_{j=1}^{n}\varrho(\lambda-q_j^-)(s_j-s_{j-1}).\quad(4.20)$$

The extremes in (4.20) are Riemann sums of the integral $\int\limits_0^t\varrho(\lambda-q(s))\,ds$; thus refining the division and observing that $\varrho(\lambda-q(s))=\varrho(\lambda)=\text{const}$ for all $\lambda\in\Omega$ if $s>t$, we obtain

$$\lim_{\tau\to\infty}\frac{\theta(t\tau,E_2)-\theta(t\tau,E_1)}{\tau\pi}=\int_0^\infty(\varrho(E_2-q(s))-\varrho(E_1-q(s)))\,ds,$$

and the assertion follows. ■

To complete the proof of Corollary 1.6, we observe that Theorem 1.2 (iii) implies that $\#_{abs}(H_\tau,\Omega^{(2)})\geqslant 1+\#_{bs}(H_\tau,\Omega)$, which together with (4.19) yields (1.19).

Acknowledgments. Various parts of this paper were written at Université de Genève, Section de Mathematiques and Mathematical Institute of the Tsukuba Univ., Japan. E. K. is grateful to these Institutes for their hospitality; he would also like to thank A. Sobolev (London) for useful discussions about the asymptotics associated with Theorem 1.6. He was supported by the 14.740.11.0581 Federal Program "Development of Scientific Potential of Higher Education for the period 2009–2013". Both authors thank the referee for helpful detailed comments.

References

[BZ] Bindel, D.; Zworski, M. Symmetry of Bound and Antibound States in the Semiclassical Limit, Letters in Mathematical Physics, 81(2007), 107-117.

[BKW] Brown, B.; Knowles, I.; Weikard, R. On the inverse resonance problem, J. London Math. Soc. (2) 68 (2003), no. 2, 383–401.

[BW] Brown, B.; Weikard, R. The inverse resonance problem for perturbations of algebro-geometric potentials. Inverse Problems 20 (2004), no. 2, 481–494.

[Ch] Christiansen, T. Resonances for steplike potentials: forward and inverse results. Trans. Amer. Math. Soc. 358 (2006), no. 5, 2071–2089.

[CL] Coddington, E.; Levinson, N. Theory of ordinary differential equations. New York, Toronto, London: McGraw-Hill 1955.

[D] Dimassi, M. Spectral shift function and resonances for slowly varying perturbations of periodic Schrödinger operators. J. Funct. Anal. 225 (2005), no. 1, 193–228.

[DG] Dyatlov, S.; Ghosh, S. Symmetry of bound and antibound states in the semiclassical limit for a general class of potentials, Proceeding of AMS, 138(2010), No 9, 3203–3210.

[E1] Eastham, M.S.P. Theory of ordinary differential equations. Van Nostrand Reinhold, London, 1970.

[E2] Eastham, M.S.P. The spectral theory of periodic differential equations. Scottish Academic Press, Edinburgh, 1973.

[F1] Firsova, N. Resonances of the perturbed Hill operator with exponentially decreasing extrinsic potential. Mat. Zametki 36 (1984), 711–724.

[F2] Firsova, N. The Levinson formula for a perturbed Hill operator. (Russian) Teoret. Mat. Fiz. 62 (1985), no. 2, 196–209.

[F3] Firsova, N. A direct and inverse scattering problem for a one-dimensional perturbed Hill operator. Mat. Sb. 130(172) (1986), no. 3, 349–385.

[F4] Firsova, N. A trace formula for a perturbed one-dimensional Schrödinger operator with a periodic potential. I. (Russian) Problems in mathematical physics, No. 7 (Russian), pp. 162–177, Izdat. Leningrad. Univ., Leningrad, 1974.

[Fr] Froese, R. Asymptotic distribution of resonances in one dimension, J. Diff. Eq., 137(1997), 251-272.

[GT] Garnett, J.; Trubowitz, E. Gaps and bands of one dimensional periodic Schrödinger operators II. Comment. Math. Helv. 62(1987), 18-37.

[G] Gérard, C. Resonance theory for periodic Schrödinger operators, Bulletin de la S.M.F., 118(1990), n0 1, 27-54.

[GS] Gesztesy, F.; Simon, B. A short proof of Zheludev's theorem. Trans. Amer. Math. Soc. 335 (1993), no. 1, 329–340.

[HKS] Hinton,D. ; Klaus, M.; Shaw, J. On the Titchmarsh-Weyl function for the half-line perturbed periodic Hill's equation. Quart. J. Math. Oxford Ser. (2) 41 (1990), no. 162, 189–224.

[H] Hitrik, M. Bounds on scattering poles in one dimension. Comm. Math. Phys. 208 (1999), no. 2, 381–411.

[Ho] Hochstadt, H. On the determination of a Hill's equation from its spectrum. Arc. Rat. Mech. Anal. 19 (1965) 353–362

[IM] Its, A.; Matveev, V. Schrödinger operators with the finite-gap spectrum and the N-soliton solutions of the Korteweg–de Vries equation. Teoret. Math. Phys. 23 (1975), 51–68.

[J] Javrjan, V. Some perturbations of self-adjoint operators. (Russian) Akad. Nauk Armjan. SSR Dokl. 38 (1964), 3–7.

[KK] Kargaev, P.; Korotyaev, E. Effective masses and conformal mappings. Comm. Math. Phys. 169 (1995), no. 3, 597–625.

[KK1] Kargaev, P.; Korotyaev, E. Inverse Problem for the Hill Operator, the Direct Approach. Invent. Math., 129 (1997), no. 3, 567–593.

[KM] Klopp, F.; Marx, M. The width of resonances for slowly varying perturbations of one-dimensional periodic Schrödingers operators, Seminaire: Equations aux Derivées Partielles. 2005–2006, Exp. No. IV, 18 pp., Semin. Equ. Deriv. Partielles, Ecole Polytech., Palaiseau.

[K1] Korotyaev, E. Inverse resonance scattering on the half line. Asymptotic Anal. 37 (2004), No 3/4, 215–226.

[K2] Korotyaev, E. Inverse resonance scattering on the real line. Inverse Problems 21 (2005), no. 1, 325–341.

[K3] Korotyaev, E. Stability for inverse resonance problem. Int. Math. Res. Not. 2004, no. 73, 3927–3936.

[K4] Korotyaev, E. Resonance theory for perturbed Hill operator. Asymptotic Anal. (to appear)

[K5] Korotyaev, E. Inverse problem and the trace formula for the Hill operator. II Math. Z. 231 (1999), no. 2, 345–368.

[K6] Korotyaev, E. Characterization of the spectrum of Schrödinger operators with periodic distributions. Int. Math. Res. Not. 2003, no. 37, 2019–2031.

[K7] Korotyaev, E. Estimates of periodic potentials in terms of gap lengths. Comm. Math. Phys. 197 (1998), no. 3, 521–526.

[Koo] Koosis, P. The logarithmic integral I, Cambridge Univ. Press, Cambridge, London, New York 1988.

[M] Marchenko, V. Sturm-Liouville operator and applications. Basel: Birkhäuser 1986.

[MO] Marchenko, V.; Ostrovski I. A characterization of the spectrum of the Hill operator. Math. USSR Sbornik 26(1975), 493-554.

[N-Z] Novikov, S.; Manakov, S. V.; Pitaevskii, L. P.; Zakharov, V. E. Theory of solitons. The inverse scattering method. Translated from the Russian. Contemporary Soviet Mathematics. Consultants Bureau [Plenum], New York, 1984.

[PT] Pöschel, P.; Trubowitz, E. Inverse Spectral Theory. Boston: Academic Press, 1987.

[Rb] Rofe-Beketov, F. A finiteness test for the number of discrete levels which can be introduced into the gaps of the continuous spectrum by perturbations of a periodic potential. Dokl. Akad. Nauk SSSR 156 (1964), 515–518.

[Sc] Schmidt, K. M. Eigenvalue asymptotics of perturbed periodic Dirac systems in the slow-decay limit. Proc. Amer. Math. Soc. 131 (2003) 1205-1214.

[S1] Simon, B. On the genericity of nonvanishing instability intervals in Hill's equation. Ann. Inst. Henri Poincaré 24 (1976), 91–93

[S2] Simon, B. Resonances in one dimension and Fredholm determinants. J. Funct. Anal. 178 (2000), no. 2, 396–420.

[So] Sobolev, A.V. Weyl asymptotics for the discrete spectrum of the perturbed Hill operator. Estimates and asymptotics for discrete spectra of integral and differential equations (Leningrad, 1989–90), 159–178, Adv. Soviet Math., 7, Amer. Math. Soc., Providence, RI, 1991.

[T] Titchmarsh, E. Eigenfunction expansions associated with second-order differential equations 2, Clarendon Press, Oxford, 1958.

[Tr] Trubowitz, E. The inverse problem for periodic potentials. Commun. Pure Appl. Math. 30(1977), 321-337.

[W] Weidmann, J. Spectral theory of ordinary differential operators. Lect. Notes in Math. 1258, Springer, Berlin, New York 1987

[Z] Zworski, M. Distribution of poles for scattering on the real line, J. Funct. Anal. 73(1987), 277-296.

[Z1] Zworski, M. SIAM, J. Math. Analysis, "A remark on isopolar potentials" 82(6), 2002, 1823-1826.

[Z2] Zworski, M. Counting scattering poles. In: Spectral and scattering theory (Sanda, 1992), 301–331, Lecture Notes in Pure and Appl. Math., 161, Dekker, New York, 1994.

[Zh1] Zheludev, V. A. The eigenvalues of a perturbed Schrödinger operator with periodic potential. (Russian) 1967 Problems of Mathematical Physics, No. 2, Spectral Theory, Diffraction Problems pp. 108–123.

[Zh2] Zheludev, V. A. The perturbation of the spectrum of the Schrödinger operator with a complex-valued periodic potential. (Russian) Problems of mathematical physics, Spectral theory 3(1968), 31–48.

[Zh3] Zheludev, V. The spectrum of Schrödinger's operator, with a periodic potential, defined on the half-axis. Works of Dept. of Math. Analysis of Kaliningrad State University (1969) (Russian), pp. 18–37.